

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 September 2001 (13.09.2001)

PCT

(10) International Publication Number
WO 01/67097 A2

(51) International Patent Classification⁷: **G01N 33/48**

(21) International Application Number: **PCT/GB01/01059**

(22) International Filing Date: **9 March 2001 (09.03.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
0005895.8 **10 March 2000 (10.03.2000)** **GB**

(71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HALE, Richard, Stephen** [GB/GB]; GlaxoSmithKline plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **ROWLEY, Adele** [GB/GB]; GlaxoSmithKline plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **BLACKSTOCK, Walter** [GB/GB]; GlaxoSmithKline plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB).

(74) Agents: **STOTT, Michael, John et al.**; GlaxoSmithKline plc, Corporate Intellectual Property, 2 New Horizons Court, Great West Road, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **NOVEL ASSAY**

(57) Abstract: A method of identifying a modulator of presenilin function, the method comprising: (i) providing (a) a polypeptide capable of binding a presenilin, which polypeptide comprises the amino acid sequence of SEQ ID NO: 1 or a functional variant thereof, or a fragment of either thereof which is capable of binding to presenilin; (b) a presenilin or a variant thereof or a fragment of either thereof capable of binding to a polypeptide which comprises the amino acid sequence of SEQ ID NO: 1; (c) a test substance under conditions that would permit binding of a polypeptide (a) to a presenilin (b) in the absence of the test substance; (ii) monitoring presenilin mediated activity; and (iii) determining thereby whether the test substance is a modulator of presenilin activity, a modulator identified by a method of the invention and use of a modulator identified by a method of the invention in the treatment of Alzheimer's disease.

BEST AVAILABLE COPY

NOVEL ASSAYField of the Invention

This invention relates to methods of identifying modulators of presenilin or catenin p120 or variants thereof and their use in the treatment of conditions in which abnormal activity of presenilin or catenin p120 is implicated such as Alzheimer's disease.

Background to the Invention

Presenilin 1 and the closely related presenilin 2 (PS1 and PS2) are membrane proteins predicted to span the membrane eight times. Presenilins are ubiquitously expressed at low levels and are located within the cell primarily in the endoplasmic reticulum (ER) and the Golgi apparatus, although presenilins have been localized to the plasma membrane in neuronal cells and COS-7 cells as well as to neuronal large dense-core granules and clathrin coated vesicles. Co-localization of PS-1 with kinetochores on the inner nuclear membrane has also been observed. Most cases of early onset familial Alzheimer's Disease (AD) are caused by mutations in the presenilin I gene. Presenilins play a role in the processing of amyloid precursor protein (APP) which is also implicated in early onset AD (reviewed by Selkoe, 1998). Co-immunoprecipitation experiments have shown that PS1 and PS2 interact directly with the immature forms of APP in the ER where the disease-associated amyloid β 1-42 peptide ($A\beta$ 42) is probably generated (Xia *et al.*, 1997; Weidemann *et al.*, 1997). Links between presenilin function and the generation of $A\beta$ 42 via intra-membrane cleavage processing of APP have been clearly demonstrated in transgenic mouse models. Presenilins may play a role in the cleavage of APP through interaction with the as yet unidentified protease termed γ -secretase or, as has recently been suggested presenilins could be the γ -secretase themselves (reviewed in Haass and Mandelkov, 1999).

Presenilin is also involved in other biological pathways. One report suggests that PS1 directly binds tau and a tau kinase, glycogen synthase kinase 3beta (GSK-3beta) and proposes that the increased association of GSK-3beta with mutant PS1 leads to increased phosphorylation of tau (Takashima *et al.* 1998).

Presenilins are also known to play an important role in Notch signalling during early embryonic development and/or cellular differentiation (reviewed by Anderton, 1999). At least one recent report claims a direct interaction between Notch1 and PS1. These results might suggest that the genetic relationship between presenilins and the Notch signalling pathway derives from a direct physical association between these proteins in the secretory pathway (Ray *et al.*, 1999).

Presenilins have also been demonstrated to interact with members of the armadillo protein family which are characterised by a series of 42 amino acid imperfect repeats that have been implicated in protein-protein interactions.

Yeast 2 hybrid experiments and/or immunoprecipitation experiments have previously shown that widely expressed β -catenins and plakoglobin/ γ -catenins and another armadillo-protein, p0071 interact with PS1 (e.g. Kang *et al.*, 1999; Stahl *et al.*, 1999; Zhang *et al.*, 1998; Zhou *et al.*, 1997b; Yu *et al.*, 1998). A neuronal-specific armadillo protein-neural plakophilin-related armadillo protein (NPRAP, a longer version of a sequence previously termed δ -catenin [Zhou *et al.*, 1997a]) also interacts with PS1 (Levesque *et al.* (1999)). Not all of these interactions are supported by all studies, for example Levesque *et al.* (1999) did not find that plakoglobin/ γ -catenin interacts with presenilin.

It has also been reported that loop regions of PS1 and PS2 interact with nonmuscle filamin (actin-binding protein 280, ABP280) and a structurally related protein (filamin homolog 1, Fh1) (Zhang *et al.*, 1998).

Catenins were discovered as proteins that are linked to the cytoplasmic domain of transmembrane cadherins. Among these junctional plaque proteins are several members of the Armadillo gene family: β -catenin, plakoglobin, and catenin p120 (p120ctn). The human p120ctn gene comprises 21 exons, potentially encoding up to 32 protein isoforms as products of alternative splicing (Keirsebilck *et al.*, 1998). Human isoforms, designated 1 to 4, differ from each other by the start codon used. Additional isoforms are derived from combinations with alternatively used exons A (exon 18) and B (20), near the end of the open reading frame, and also with exon C (11) in the middle of the open reading frame. Hence, the longest isoform is of type 1ABC and comprises 968 amino acid residues. The functional consequence of the observed multitude of p120ctn splice variants is unclear but there are tissue-specific

expression level differences. The exon organization, which is not simply related to the Armadillo repeat structure, is very well conserved between the p120ctn gene and the related ARVCF gene, but not at all between these two genes and the β -catenin or plakoglobin genes.

Summary of the Invention

The present invention is based on a novel direct action between a presenilin and a protein, catenin p120.

The DNA and predicted amino acid sequence of catenin p120 have been previously deposited in public domain databases but this gene product has not previously been shown to interact with presenilin.

The interaction between presenilin and catenin p120 provides a new therapeutic intervention point in disorders involving defective presenilin function, and more specifically in Alzheimer's disease. In addition catenin p120 is now proposed as a target for identifying agents which may be useful in the treatment of Alzheimer's disease.

Accordingly the invention provides:
a method of identifying a modulator of presenilin function, the method comprising:

(i) providing

- (a) a polypeptide capable of binding a presenilin, which polypeptide comprises the amino acid sequence of SEQ ID NO: 1 or a functional variant thereof or a functional fragment of either thereof which is capable of binding to a presenilin;
- (b) a presenilin or a variant thereof or a fragment of either thereof capable of binding to a polypeptide which comprises the amino acid sequence of SEQ ID NO: 1; and
- (c) a test substance

under conditions that would permit binding of (a) to (b) in the absence of (c);

- (ii) monitoring presenilin mediated activity; and
- (iii) determining thereby whether the test substance is a modulator of presenilin activity.

In a further aspect, the invention provides a method for identification of a compound that modulates catenin p120 activity, which method comprises:

- (i) contacting a catenin p120 polypeptide comprising
 - (a) the amino acid sequence of SEQ ID NO: 1; or
 - (b) a variant thereof or a fragment of either thereof which maintains a catenin p120 function; with a test compound and
- (ii) monitoring for catenin p120 activity thereby determining whether the test compound is a modulator of catenin p120.

The invention also provides:

- a modulator identified by a method according to any one of the preceding claims for use in a method of treatment of the human or animal body by therapy;
- use of a modulator identified by a method according to any one of claims 1 to 10 in the manufacture of a medicament for the treatment or prophylaxis of a disorder that is responsive to modulation of presenilin activity;
- use of a polypeptide capable of binding a presenilin, which polypeptide comprises the amino acid sequence of SEQ ID NO: 1 or a variant or fragment of SEQ ID NO: 1 which is capable of binding a presenilin in the manufacture of a medicament for the treatment, prophylaxis or diagnosis of a disorder that is responsive to modulation of presenilin activity;
- use of a polynucleotide which encodes a polypeptide as defined in claim 13 or a catenin p120 polypeptide as defined in claim 9 comprising:

- (a) the sequence of SEQ ID NO: 2; or
- (b) a sequence that hybridizes to the complement of SEQ ID NO: 2; or
- (c) a sequence that is degenerate as a result of the genetic code with respect to a sequence defined in (a) or (b); or
- (d) a sequence that is complementary to a polynucleotide defined in (a), (b) or (c);

in the manufacture of a medicament for the treatment, prophylaxis or diagnosis of a disorder that is responsive to modulation of presenilin activity.

Preferably the disorder responsive to presenilin modulation is Alzheimer's disease.

Brief Description of the Sequences

5 SEQ ID NO: 1 is the amino acid sequence of catenin p120.

SEQ ID NO: 2 comprises the amino acid sequence and encoding polynucleotide of catenin p120.

SEQ ID NO: 3 is the amino acid sequence of presenilin 1.

SEQ ID NO: 4 is the amino acid and encoding polynucleotide sequence of presenilin

10 1.

SEQ ID NO: 5 is the amino acid sequence of presenilin 2.

SEQ ID NO: 6 is the amino acid and encoding polynucleotide sequence of presenilin 2.

Detailed Description of the Invention

The invention provides a method for identifying a modulator of presenilin activity or a modulator of catenin p120 activity. A modulator may modulate the interaction between a presenilin and a catenin p120 isoform.

20 A presenilin for use in accordance with the invention herein referred to as presenilin (b) may comprise a naturally occurring presenilin such as PS-1 or PS-2 having the amino acid sequence of SEQ ID No: 3 or SEQ ID NO: 5 or may comprise a variant or fragment of such naturally occurring presenilin such as another unidentified isoform or splice variant which is homologous to or retains the desired function of a known presenilin. Such a variant or fragment of presenilin for use in
25 the invention is one which is capable of binding to catenin p120 having the sequence of SEQ ID NO: 1.

A polypeptide for use in accordance with the invention is one which capable of binding presenilin. The polypeptide, herein referred to as polypeptide (a) a
30 catenin p120 isoform having the sequence of SEQ ID NO: 1 or a functional variant or a functional fragment thereof. A variant may comprise a naturally occurring isoform or splice variant. A variant or fragment of SEQ ID NO: 1 for use in accordance with the invention is capable of binding to presenilin and in particular

presenilin 1.

To determine whether a variant or fragment of SEQ ID NO: 1 is capable of binding to a presenilin the variant or fragment can be contacted with a presenilin under conditions suitable for the formation of a complex between catenin p120 and presenilin. Similarly, to determine whether a presenilin, or variant or fragment thereof is capable of binding to catenin p120, the presenilin or fragment thereof can be contacted with catenin p120 under conditions suitable for the formation of a complex between catenin p120 and presenilin. Any one of the assays described herein can be carried out in the absence of a test substance to determine the binding capabilities of these proteins.

Proteins with naturally occurring amino acid sequences are preferred for use in the assays. Preferred proteins are human proteins but homologues from other mammalian species, or other animal species may be used. Any allelic variant or species homologue of the defined proteins may be used. References to a variant of the protein as described below relates to a variant of a presenilin or catenin p120. For all the proteins described herein for use in an assay of the invention, the ability of the variant to bind catenin p120 or presenilin as appropriate is preferably maintained.

Allelic variants and species homologues can be obtained by following the procedures described herein for the identification and production of proteins that bind catenin p120 or a presenilin as appropriate. It is also possible to use a nucleic acid probe as described herein to probe libraries made from human or other animal cells in order to obtain clones encoding allelic or species variants. The clones can be manipulated by conventional techniques to generate a polypeptide which can then be produced by recombinant or synthetic techniques known in the art.

Polypeptides that have been artificially mutated but retain catenin p120 or presenilin binding activity or other catenin p120 or presenilin activity may also be used in the invention. Such mutants may be generated by techniques well known in the art, including site directed mutagenesis, random mutagenesis and restriction enzyme digestion and ligation. A protein for use in the invention preferably has at least 60% sequence identity to a natural protein, more preferably at least 70%, at least 80%, at least 90%, at least 95%, at least 97% or at least 99% sequence identity thereto over a region of at least 20, preferably at least 30, for instance at least 40, at

least 60, at least 100 contiguous amino acids or over the full length of a natural protein. Amino acid substitutions may be made, for example from 1, 2 or 3 to 10, 20 or 30 substitutions. Conservative substitutions may be made, for example according to the following Table. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other.

ALIPHATIC	Non-polar	G A P
		I L V
	Polar-uncharged	C S T M
		N Q
	Polar-charged	D E
		K R
AROMATIC		H F W Y

The entire protein sequence of each of the proteins used in the assay may be present. Fragments of the proteins and variants described above that retain the ability to bind to the second component in the binding assay, i.e. presenilin for catenin p120 and catenin p120 for presenilin may also be used in the invention. Alternatively variants or fragments of catenin p120 which retain a function of catenin p120 may be used in assays to identify modulators of catenin p120. Preferred fragments will be at least 30, e.g. at least 100, at least 200, at least 300, at least 400, at least 500 or at least 600 amino acids in size. Preferred presenilin fragments are stable N- and C- terminal fragments that are generated in vivo by endoproteolysis (Capell *et al.*, 1998; Yu *et al.*, 1998). The term "isoform" is used herein to describe such variants and fragments of catenin p120 or presenilin.

The polypeptides for use in the invention may be chemically modified, e.g. post-translationally modified. For example, they may be glycosylated or comprise modified amino acid residues. They may also be modified by the addition of a signal sequence to promote their secretion from a cell where the polypeptide does not naturally contain such a sequence. The polypeptides may be tagged to aid detection or purification, for example using a HA, his8, his6, T7, myc or flag tag.

Alternatively the polypeptides may be fusion proteins to aid purification or detection, for example, GST-fusion proteins may be used to aid purification from bacteria and GFP-fusion proteins may be used to aid detection. The polypeptide (a) and presenilin (b) may be tagged with different labels which may assist in identification of a polypeptide/presenilin complex.

Polypeptides for use in the invention may be in a substantially isolated form. It will be understood that the polypeptide may be mixed with carriers or diluents which will not interfere with the intended purpose of the polypeptide and still be regarded as substantially isolated. A polypeptide for use in the invention may also be in a substantially purified form, in which case it will generally comprise the polypeptide in a preparation in which more than 50%, e.g. more than 80%, 90%, 95% or 99%, by weight of the polypeptide in the preparation is a polypeptide of the invention.

Polynucleotides

The invention provides a polynucleotide which encodes a polypeptide capable of binding a presenilin which polynucleotide consists essentially of:

- (a) the sequence of SEQ ID NO: 2; or
- (b) a sequence that hybridizes to the complement of SEQ ID NO: 2; or
- (c) a sequence that is degenerate as a result of the genetic code with respect to a sequence defined in (a) or (b); or
- (d) a sequence that is complementary to a polynucleotide defined in (a), (b) or (c);

for use in a method of treatment of the human or animal body by therapy.

Typically a polynucleotide of the invention comprises a contiguous sequence of nucleotides which is capable of hybridizing under selective conditions to the complement of the coding sequence of SEQ ID NO: 2. Preferably a polynucleotide of the invention encodes a polypeptide which is capable of binding to a presenilin or retains a function of catenin p120.

A polynucleotide comprising a sequence that hybridizes to the complement of the coding sequence of SEQ ID NO: 2 can hybridize at a level significantly above background. Background hybridization may occur, for example, because of other cDNAs present in a cDNA library. The signal level generated by the interaction

between a polynucleotide of the invention and the complement of the coding sequence of SEQ ID NO: 2 is typically at least 10 fold, preferably at least 100 fold, as intense as interactions between other polynucleotides and the coding sequence of SEQ ID NO: 2. The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ^{32}P . Selective hybridisation may typically be achieved using conditions of low stringency (0.3M sodium chloride and 0.03M sodium citrate at about 40°C), medium stringency (for example, 0.3M sodium chloride and 0.03M sodium citrate at about 50°C) or high stringency (for example, 0.03M sodium chloride and 0.003M sodium citrate at about 60°C). However, such hybridization may be carried out under any suitable conditions known in the art (see Sambrook *et al.*, 1989). For example, if high stringency is required, suitable conditions include 0.2 X SSC at 60°C. If lower stringency is required, suitable conditions include 2 X SSC at 60°C.

A nucleotide sequence which is capable of selectively hybridizing to the complement of the DNA coding sequence of SEQ ID NO: 2 will generally have at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 98% or at least 99% sequence identity to the coding sequence of SEQ ID NO: 2 over a region of at least 20, preferably at least 30, for instance at least 40, at least 60, more preferably at least 100 contiguous nucleotides or most preferably over the full length of SEQ ID NO: 2. Methods of measuring nucleic acid and protein homology are well known in the art.

For example the UWGCG Package provides the BESTFIT program which can be used to calculate homology (for example used on its default settings) (Devereux *et al* (1984) *Nucleic Acids Research* 12, p387-395). The PILEUP and BLAST algorithms can be used to calculate homology or line up sequences (typically on their default settings), for example as described in Altschul (1993) *J. Mol. Evol.* 36:290-300; Altschul *et al* (1990) *J. Mol. Biol.* 215:403-10.

Software for performing BLAST analyses is publicly available through the National Centre for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>).

This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length

in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al.*, 1990). These initial neighbourhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90: 5873-5787. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

Any combination of the above mentioned degrees of sequence identity and minimum sizes may be used to define polynucleotides for use in the invention, with the more stringent combinations (i.e. higher sequence identity over longer lengths) being preferred. Thus, for example a polynucleotide which has at least 90% sequence identity over 25, preferably over 30 nucleotides forms one aspect of the invention, as does a polynucleotide which has at least 95% sequence identity over 40 nucleotides.

The coding sequence of SEQ ID NO: 2 may be modified by nucleotide substitutions, for example from 1, 2 or 3 to 10, 25, 50 or 100 substitutions. The polynucleotide of SEQ ID NO: 2 may alternatively or additionally be modified by one or more insertions and/or deletions and/or by an extension at either or both ends.

The modified polynucleotide generally encodes a protein that can bind a presenilin. Degenerate substitutions may be made and/or substitutions may be made which would result in a conservative amino acid substitution when the modified sequence is translated, for example as shown in the Table above.

5 Polynucleotides of the invention may comprise DNA or RNA. They may also be polynucleotides which include within them synthetic or modified nucleotides. A number of different types of modification to polynucleotides are known in the art. These include methylphosphonate and phosphorothioate backbones, addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the
10 purposes of the present invention, it is to be understood that the polynucleotides described herein may be modified by any method available in the art. Such modifications may be carried out in order to enhance the *in vivo* activity or lifespan of polynucleotides of the invention.

Polynucleotides according to the invention may be produced recombinantly,
15 synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques. The polynucleotides are typically provided in isolated and/or purified form.

Although in general the techniques mentioned herein are well known in the art, reference may be made in particular to Sambrook *et al*, 1989, Molecular Cloning:
20 a laboratory manual.

A polynucleotide may be an essential component in an assay of the invention, a probe (or template for designing a probe) for identifying proteins that may be used in the invention or a test compound. The nucleotides according to the invention have utility in production of the proteins according to the invention, which may take place
25 *in vitro*, *in vivo* or *ex vivo*. The nucleotides may be involved in recombinant protein synthesis or indeed as therapeutic agents in their own right, utilised in gene therapy techniques. Antisense sequences, may also be used in gene therapy, such as in strategies for down regulation of expression of the proteins of the invention. The invention further provides double stranded polynucleotides comprising a
30 polynucleotide for use in the invention and its complement for use in a method of treatment of the human or animal body by therapy.

Probes and other fragments will preferably be at least 10, preferably at least

15 or at least 20, for example at least 25, at least 30 or at least 40 nucleotides in length. They will typically be up to 40, 50, 60, 70, 100 or 150 nucleotides in length. Probes and fragments can be longer than 150 nucleotides in length, for example up to 200, 300, 400, 500, 600, 700 nucleotides in length, or even up to a few nucleotides, such as five or ten nucleotides, short of the coding sequence of SEQ ID NO: 2.

Polynucleotides of the invention or for use in the invention can be inserted into expression vectors. Such expression vectors are routinely constructed in the art of molecular biology and may for example involve the use of plasmid DNA and appropriate initiators, promoters, enhancers and other elements, such as for example polyadenylation signals which may be necessary, and which are positioned in the correct orientation, in order to allow for protein expression. Other suitable vectors would be apparent to persons skilled in the art. By way of further example in this regard we refer to Sambrook *et al.*

Polynucleotides according to the invention may also be inserted into the vectors described above in an antisense orientation in order to provide for the production of antisense RNA. Antisense RNA or other antisense polynucleotides may also be produced by synthetic means. Such antisense polynucleotides may be used as test compounds in the assays of the invention or may be useful in a method of treatment of the human or animal body by therapy.

Preferably, a polynucleotide of the invention or for use in the invention in a vector is operably linked to a control sequence which is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A regulatory sequence, such as a promoter, "operably linked" to a coding sequence is positioned in such a way that expression of the coding sequence is achieved under conditions compatible with the regulatory sequence.

Vectors for use in the invention may be transformed into a suitable host cell as described above to provide for expression of a polypeptide for use in the invention. Thus, in a further aspect the invention provides a process for preparing a polypeptide according to the invention which comprises cultivating a host cell transformed or transfected with an expression vector encoding the polypeptide, and

recovering the expressed polypeptide.

The vectors may be for example, plasmid, virus or phage vectors provided with a origin of replication, optionally a promoter for the expression of the said polynucleotide and optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a resistance gene for a fungal vector. Vectors may be used *in vitro*, for example for the production of RNA or used to transfect or transform a host cell, for example, a mammalian host cell. The vectors may also be adapted to be used *in vivo*, for example in a method of gene therapy.

Promoters and other expression regulation signals may be selected to be compatible with the host cell for which expression is designed. For example, yeast promoters include *S. cerevisiae* GAL4 and ADH promoters, *S. pombe* *nmt1* and *adh* promoter. Mammalian promoters include the metallothionein promoter which can be induced in response to heavy metals such as cadmium. Viral promoters such as the SV40 large T antigen promoter or adenovirus promoters may also be used. All these promoters are readily available in the art.

Mammalian promoters, such as β -actin promoters, may be used. Tissue-specific promoters, in particular endothelial or neuronal cell specific promoters (for example the DDAHI and DDAHII promoters), are especially preferred. Viral promoters may also be used, for example the Moloney murine leukaemia virus long terminal repeat (MMLV LTR), the rous sarcoma virus (RSV) LTR promoter, the SV40 promoter, the human cytomegalovirus (CMV) IE promoter, adenovirus, HSV promoters (such as the HSV IE promoters), or HPV promoters, particularly the HPV upstream regulatory region (URR). Viral promoters are readily available in the art.

The vector may further include sequences flanking the polynucleotide which comprise sequences homologous to eukaryotic genomic sequences, preferably mammalian genomic sequences, or viral genomic sequences. This will allow the introduction of the polynucleotides of the invention into the genome of eukaryotic cells or viruses by homologous recombination. In particular, a plasmid vector comprising the expression cassette flanked by viral sequences can be used to prepare a viral vector suitable for delivering the polynucleotides of the invention to a mammalian cell. Other examples of suitable viral vectors include herpes simplex

viral vectors and retroviruses, including lentiviruses, adenoviruses, adeno-associated viruses and HPV viruses (such as HPV-16 or HPV-18). Gene transfer techniques using these viruses are known to those skilled in the art. Retrovirus vectors for example may be used to stably integrate the polynucleotide giving rise to the antisense RNA into the host genome. Replication-defective adenovirus vectors by contrast remain episomal and therefore allow transient expression.

The invention also utilizes cells that have been modified to express a presenilin polypeptide and/or a catenin p120 polypeptide. Such cells include transient, or preferably stable higher eukaryotic cell lines, such as mammalian cells or insect cells, using for example a baculovirus expression system, lower eukaryotic cells, such as yeast or prokaryotic cells such as bacterial cells. Particular examples of cells which may be modified by insertion of vectors encoding for a polypeptide according to the invention include mammalian HEK293T, CHO, HeLa and COS cells. Preferably the cell line selected will be one which is not only stable, but also allows for mature glycosylation of a polypeptide. A polypeptide of the invention may be overexpressed in bacterial cells, such as *E.coli*, and isolated from the bacterial culture.

According to another aspect, the present invention may also use antibodies (either polyclonal or preferably monoclonal antibodies, chimeric, single chain, Fab fragments) which have been raised by standard techniques and are specific for a presenilin polypeptide or for a catenin p120 polypeptide. Such antibodies could for example, be useful in purification, isolation or screening methods involving immunoprecipitation techniques and may be used as tools to further elucidate the function of catenin p120 or a variant thereof, or indeed as therapeutic agents in their own right. Antibodies may also be raised against specific epitopes of the proteins according to the invention. Such antibodies may be used to block ligand binding to the receptor. An antibody, or other compounds, "specifically binds" to a protein when it binds with preferential or high affinity to the protein for which it is specific but does not bind or binds with only low affinity to other proteins. A variety of protocols for competitive binding or immunoradiometric assays to determine the specific binding capability of an antibody are well known in the art (see for example Maddox *et al* 1993). Such immunoassays typically involve the formation of

complexes between the "specific protein" and its antibody and the measurement of complex formation.

Antibodies may be antibodies to human polypeptides or fragments thereof. For the purposes of this invention, the term "antibody", unless specified to the contrary, includes fragments which bind a polypeptide of the invention. Such fragments include Fv, F(ab') and F(ab')₂ fragments, as well as single chain antibodies. Furthermore, the antibodies and fragment thereof may be chimeric antibodies, CDR-grafted antibodies or humanised antibodies.

Antibodies can be produced by any suitable method. Means for preparing and characterising antibodies are well known in the art, see for example Harlow and Lane (1988) "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. For example, an antibody may be produced by raising antibody in a host animal against the whole polypeptide or a fragment thereof, for example an antigenic epitope thereof, herein after the "immunogen".

A method for producing a polyclonal antibody comprises immunising a suitable host animal, for example an experimental animal, with the immunogen and isolating immunoglobulins from the animal's serum. The animal may therefore be inoculated with the immunogen, blood subsequently removed from the animal and the IgG fraction purified.

A method for producing a monoclonal antibody comprises immortalising cells which produce the desired antibody. Hybridoma cells may be produced by fusing spleen cells from an inoculated experimental animal with tumour cells (Kohler and Milstein (1975) *Nature* 256, 495-497).

An immortalized cell producing the desired antibody may be selected by a conventional procedure. The hybridomas may be grown in culture or injected intraperitoneally for formation of ascites fluid or into the blood stream of an allogenic host or immunocompromised host. Human antibody may be prepared by *in vitro* immunisation of human lymphocytes, followed by transformation of the lymphocytes with Epstein-Barr virus.

For the production of both monoclonal and polyclonal antibodies, the experimental animal is suitably a goat, rabbit, rat or mouse. If desired, the immunogen may be administered as a conjugate in which the immunogen is coupled,

for example via a side chain of one of the amino acid residues, to a suitable carrier. The carrier molecule is typically a physiologically acceptable carrier. The antibody obtained may be isolated and, if desired, purified.

Assays

5 Any suitable assay format may be used for identifying a modulator of a presenilin, for example a modulator of a presenilin/ catenin p120 interaction. In a preferred aspect, the assay is a cell-based assay. Preferably the assay may be carried out in a single well of a microtitre plate. Assay formats which allow high throughput screening are preferred.

10 As the first step of the method for identifying a modulator of presenilin function, (a) a polypeptide comprising the sequence of SEQ ID NO: 1 or a variant or fragment of either sequence capable of binding to a presenilin; (b) a presenilin or a variant or a fragment thereof capable of binding to catenin p120; and (c) a test substance are contacted under conditions that would permit binding of (a) to (b) in
15 the absence of a test substance. The activity of presenilin is then monitored. For example, the interaction between the polypeptide (a) and a presenilin (b) may be analysed. The interaction between the polypeptide(a) and presenilin (b) in the presence of a test substance may be compared with the interaction between the polypeptide (a) and presenilin (b) in the absence of the test substance to determine
20 whether the test substance modulates the binding of polypeptide (a) and presenilin (b) and thereby whether the test substance enhances or inhibits the binding of a presenilin to catenin p120.

As used herein, a polypeptide (a) is used to refer to catenin p120 having the sequence of SEQ ID NO: 1 or an isoform or variant thereof or a fragment of any
25 thereof which is capable of binding to a presenilin, or to a variant or a fragment of presenilin which is capable of binding to catenin p120.

The test substance can be contacted with a cell harbouring a polynucleotide or expression vector encoding a polypeptide (a) and a polynucleotide or expression vector encoding a presenilin (b). Optionally the cell may harbour a polynucleotide or
30 expression vector encoding a test substance, wherein the test substance is a peptide. The cell typically allows transcription and translation of the polynucleotides or vectors so that the polypeptides are expressed in the same cell.

The test substance may be provided in the extracellular medium used for washing, incubating or growing the cell. The test substance may modulate the interaction of presenilin (b) with the polypeptide (a) indirectly from outside the cell, for example by interacting with an extracellular domain of presenilin or may be taken up into the cell from the extracellular medium. Where presenilin (b) and polypeptide (a) are coexpressed in a cell, the cell may express both proteins naturally, for example the cell may be a neuronal cell grown in a primary culture, or the cell may express both proteins recombinantly, or the cell may naturally express one protein and be transformed to express the other protein recombinantly.

The cell may be transiently or stably transfected or transformed. The polypeptide (a), presenilin (b) and the test substance (c) where it is a peptide may all be stably expressed. More preferably polypeptide (a) and presenilin (b) will be stably expressed and a test substance peptide will be transiently expressed. Where only polypeptide (a) and presenilin (b) are expressed by the cell they may both be transiently expressed, both stably expressed or one may be stably expressed and the other transiently expressed. Cells can be transfected by methods well known in the art, for example, by electroporation, calcium phosphate precipitation, lipofection or heat shock. The proteins may be expressed in mammalian cells such as human cells or non-mammalian cells such as yeast or bacteria. It is preferred that the cells are in culture. Preferred cell lines which may be used include HEK293, COS and PC12 cells.

A polypeptide (a) and a presenilin (b) can also be recombinantly expressed in different cells. These cells may be two different cultures of the same cell line or may be different cell lines. The cell lines may both be mammalian cells, bacterial cells or yeast cells or the two cell lines may be from different organisms, for example a polypeptide (a) may be expressed in a mammalian cell and a presenilin (b) in a bacterial cell. A cell expressing a polypeptide (a) or a cell lysate, a membrane preparation or a protein preparation derived from a cell expressing a polypeptide (a) can be contacted with a cell expressing a presenilin (b) or a cell homogenate, a cell lysate or a protein preparation from cells expressing presenilin (b). Similarly, where the test substance is a peptide or protein, an expression vector comprising a polypeptide encoding (c) can be singly transfected into a cell and the cell

homogenate, cell lysate, membrane preparation or protein preparation from the transfected cell can be used in the assay.

The conditions which permit binding of a polypeptide (a) to a presenilin (b) in an extracellular environment can be determined by carrying out the assay in the absence of a test substance.

A number of biochemical and molecular cell biology protocols known in the art can be used to analyse the interaction of a polypeptide (a) and presenilin (b) (see for example Sambrook *et al.*, 1989). Some specific examples are outlined below:

The presenilin/catenin p120 interaction can be determined directly by incubating a radiolabelled polypeptide (a) with the presenilin and monitoring binding of the polypeptide (a) to the presenilin. Alternatively binding of radiolabelled presenilin to catenin p120 polypeptide (a) may be monitored. Typically, the radiolabelled substance can be incubated with cell membranes containing the presenilin or catenin p120 until equilibrium is reached. The membranes can then be separated from a non-bound substance and dissolved in scintillation fluid to allow the radioactive content to be determined by scintillation counting. Non-specific binding of the substance may also be determined by repeating the experiment in the presence of a saturating concentration of a non-radioactive polypeptide (a) or presenilin. Preferably a binding curve is constructed by repeating the experiment with various concentrations of the radiolabelled substance.

A yeast-2 hybrid assay system may be used. A polynucleotide encoding a presenilin, or fragment thereof capable of binding to catenin p120 can be cloned into GAL4 binding domain vector (GAL4_{BD}) and a polynucleotide comprising the sequence of SEQ ID NO: 2 or a variant or fragment thereof capable of binding to a presenilin can be cloned into a GAL4 activation domain fusion vector (GAL4_{AD}). Alternatively, a polynucleotide comprising the sequence of SEQ ID NO: 2 or a variant or fragment thereof capable of binding to a presenilin can be cloned into GAL4_{BD} and a polynucleotide encoding a presenilin can be cloned into GAL4_{AD}. GAL4_{AD} and GAL4_{BD} can then be expressed in yeast and the resulting β -galactosidase activity can be assayed and quantified using the substrate o-nitrophenol β -D-galactopyranoside (ONPG) using a liquid nitrogen freeze fracture regime as described by Harshman *et al.*, 1998.

A "pull-down" assay system may also be used. A presenilin, or a variant or fragment thereof capable of binding a catenin p120 can be run on a denaturing SDS polyacrylamide gel and transferred to a nitrocellulose membrane. Use of a fusion protein, such as a GST-fusion protein expressed and purified from *E. coli*, is preferred. The protein on the gel can then be renatured. A number of protocols for the refolding of denatured proteins are detailed in Marston (1987). In a parallel experiment, the position of the presenilin can be identified by immunoblotting using techniques well known in the art. A labelled polypeptide (a) capable of interacting with a presenilin, which may be present, for example, in a cell extract from cells transiently transfected with a polynucleotide of SEQ ID NO: 2 capable of interacting with a presenilin grown in medium containing ³⁵S-methionine, can then be incubated with the nitrocellulose membrane. The test compound can be included in the incubation medium. After washing to remove non-specifically bound proteins, labelled catenin p120 bound to the presenilin can be detected and quantified using, for example, a phosphorimager or a scintillation counter. The assay can also be carried out by immobilizing the catenin p120 and measuring the binding of a presenilin to the immobilized protein.

Alternatively, a presenilin or variant or fragment thereof capable of binding catenin p120; a polypeptide comprising SEQ ID NO: 1 or a variant or fragment thereof capable of interacting with a presenilin may be immunoprecipitated, immunopurified or affinity purified from a cell extract of cells co-expressing a presenilin or fragment thereof and a polypeptide comprising SEQ ID NO: 1 or a variant or fragment of either sequence capable of binding to a presenilin. If the test substance is a polypeptide the cells may also co-express the test substance.

Alternatively, the test substance may be provided in the cell growth medium. Coprecipitating/copurifying catenin p120 or presenilins can then be detected, for example using Western blotting techniques or by radiolabelling recombinantly expressed proteins, and quantified using a phosphorimager or scintillation counter.

An important aspect of the present invention is the use of catenin p120 polypeptides according to the invention in screening methods to identify compounds that may act as modulators of catenin p120 activity and in particular compounds that may be useful in treating presenilin associated disease. Any suitable form may be

used for the assay to identify a modulator of catenin p120 activity. In general terms, such screening methods may involve contacting a polypeptide of the invention with a test compound and then measuring activity.

Any suitable activity of catenin p120 may be measured in a screening assay.

5 For example, the binding of catenin p120 to the cytoskeleton or to a transcription factor may be monitored. A modulator may, for example, modulate binding of catenin p120 to E-cadherin and/or may modulate binding of catenin p120 to a transcription factor such as Kaiso (Daniel and Reynolds, 1999). Alternatively, catenin p120 activity may be monitored by detecting expression of a reporter gene,
10 such as luciferase, under the control of a regulatory sequence which binds a catenin p120/transcription factor protein complex or a transcription factor which is activated by catenin p120.

A ligand of catenin p120 can be determined directly by incubating a radiolabelled test substance with the polypeptide (a) and monitoring binding of the
15 test compound to the polypeptide. Typically, the radiolabelled test substance can be incubated with cell membranes containing the polypeptide until equilibrium is reached. The membranes can then be separated from a non-bound test substance and dissolved in scintillation fluid to allow the radioactive content to be determined by scintillation counting. Non-specific binding of the test substance may also be
20 determined by repeating the experiment in the presence of a saturating concentration of a non-radioactive ligand. Preferably a binding curve is constructed by repeating the experiment with various concentrations of the test substance.

Modulator activity can be determined by contacting cells expressing a polypeptide (a) of the invention with a substance under investigation and by
25 monitoring the effect mediated by the polypeptide. The cells expressing the polypeptide may be *in vitro*, for example in cultured cells, or *in vivo*. The polypeptide of the invention may be naturally or recombinantly expressed. Preferably, the assay is carried out *in vitro* using cells expressing recombinant polypeptide.

30 For each assay system a parallel control experiment (in which the substance to be tested is omitted) and experiments in which a test substance is included can be carried out. The results of the experiments using the test compound and the control

experiments can be used to determine whether the test compound inhibits or enhances binding.

The substance tested may be tested with any other known receptor/interacting cytoplasmic protein combinations, for example presenilin and β -catenin, to exclude the possibility that the test substance is a general inhibitor of protein/protein interactions.

Where a variant or fragment of presenilin or catenin p120 is used in the assay as the presenilin (b) or polypeptide (a), the assay is preferably run first in the absence of a test substance to ensure that the variant or fragment does not affect the activity of the presenilin or catenin p120. The assays may also be carried out monitoring PS mediated signalling. For example, the step of monitoring presenilin activity may involve assessment of presenilin mediated signalling or processing or the effect of binding of presenilin to other proteins. For example the assay may involve determination of APP processing, Notch signalling, for example in early development, or the binding of presenilin to tau and a tau kinase.

Candidate Modulators

A modulator of presenilin or catenin p120 function may exert its effect by binding directly to presenilin or catenin p120 polypeptide or may have an upstream effect which prevents the presenilin/catenin p120 interaction occurring, or presenilin or catenin p120 mediated activity.

A modulator may directly inhibit the interaction of presenilin with catenin p120 or inhibit interaction between catenin p120 and a ligand such as E-cadherin. A candidate modulator may comprise a fragment of a catenin p120 isoform capable of binding a presenilin or catenin p120 ligand but lacking any functional activity. Alternatively, a candidate modulator may comprise a fragment of a presenilin capable of binding catenin p120 but lacking any functional activity. Candidate molecules include N- terminal and C- terminal fragments of presenilin 1 or presenilin 2. Antibodies or antibody fragments, for example as defined herein, that specifically bind to presenilin or catenin p120 or chemical compounds capable of binding these proteins are also candidate compounds.

Other suitable test substances include combinatorial libraries, defined chemical entities and compounds, peptide and peptide mimetics, oligonucleotides

and natural product libraries, such as display libraries (e.g. phase display libraries).

Typically, organic molecules will be screened, preferably small organic molecules which have a molecular weight of from 50 to 2500 daltons. Candidate products can be biomolecules including, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs.

Test substances may be used in an initial screen of, for example, 10 substances per reaction, and the substances of these batches which show inhibition or activation tested individually. Test substances may be used at a concentration of from 1nM to 10mM, preferably from, 100nM to 1000 μ M or from 1 μ M to 100 μ M, more preferably from 1 μ M to 10 μ M.

Modulators

A modulator of presenilin activity which produces a measurable reduction or increase in catenin p120 to presenilin in the assays described above, or an effect on presenilin activity or catenin p120 mediated activity.

Preferred inhibitors are those which inhibit binding by at least 10%, at least 20%, at least 30%, at least 40% at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 99% at a concentration of the inhibitor of 1 μ g ml⁻¹, 10 μ g ml⁻¹, 100 μ g ml⁻¹, 500 μ g ml⁻¹, 1mg ml⁻¹, 10mg ml⁻¹ or 100mg ml⁻¹.

Preferred activators are those which activate binding by at least 10%, at least 25%, at least 50%, at least 100%, at least, 200%, at least 500% or at least 1000% at a concentration of the activator 1 μ g ml⁻¹, 10 μ g ml⁻¹, 100 μ g ml⁻¹, 500 μ g ml⁻¹, 1mg ml⁻¹, 10mg ml⁻¹ or 100mg ml⁻¹.

The percentage inhibition or activation represents the percentage decrease or increase in expression/activity in a comparison of assays in the presence and absence of the test substance. Any combination of the above mentioned degrees of percentage inhibition or activation and concentration of inhibitor or activator may be used to define an inhibitor or activator of the invention, with greater inhibition or activation at lower concentrations being preferred.

Candidate substances which show activity in assays such as those described above can be tested in *in vivo* systems, an animal model. Candidate inhibitors could be tested for their ability to decrease presenilin mediated signalling, for example by interfering with development in *C.Elegans* or by interfering with APP processing or by monitoring changes in gene expression using a reporter gene assay.

Candidate activators could be tested for their ability to increase presenilin mediated signalling. Ultimately such substances would be tested in animal models of the target disease states.

Therapeutic use

Modulators of presenilin/catenin p120 or of presenilin activity or of catenin p120 activity identified by the methods of the invention may be used for the treatment or prophylaxis of a disorder that is responsive to modulation of presenilin activity or catenin p120.

In particular, neuronal disorders such as cognitive disorders including Alzheimer's disease may be treated. A modulator of presenilin or catenin p120 activity may be used to alleviate the symptoms or to improve the condition of a patient suffering from such a disorder. A therapeutically effective amount of a modulator is an amount which is sufficient to alleviate one or more symptoms of a disorder or to improve the condition of a patient suffering from a disorder.

Modulators of presenilin or catenin p120 activity may be useful in enhancing cognitive function. Hence, a therapeutically effective amount of a modulator may be an amount which is sufficient to produce an enhancement of cognitive function in a patient suffering from a neurodegenerative disorder. This may be useful in treating neurodegenerative diseases such as Alzheimer's disease or in enhancing cognitive function following injury to the brain.

Catenin p120 polypeptides and polynucleotides as described herein may also be used in the treatment or prophylaxis of such disorders.

Another aspect of the present invention is the use of polynucleotides encoding the catenin p120 polypeptides of the invention to identify mutations in catenin p120 genes which may be implicated in human disorders. Identification of such mutations may be used to assist in diagnosis of or susceptibility to Alzheimer's or other conditions associated with presenilin and in assessing the physiology of such

disorders. Polynucleotides may also be used in hybridisation studies to monitor for expression of catenin p120 genes and in particular for up or down regulation of catenin p120 expression.

The present invention provides a method for assessing a disorder associated with abnormal presenilin function such as Alzheimer's disease by detecting variation in the expressed products encoded by a catenin p120 gene. This may comprise determining the level of an catenin p120 expressed in cells or determining specific alterations in the expressed product. Sequences of interest for diagnostic purposes include, but are not limited to, the conserved portions as identified by sequence similarity and conservation of intron/exon structure. The diagnosis may be performed in conjunction with kindred studies to determine whether a mutation of interest co-segregates with disease phenotype in a family.

Diagnostic procedures may be performed on polynucleotides isolated from an individual or alternatively, may be performed *in situ* directly upon tissue sections (fixed and/or frozen) of patient tissue obtained from biopsies or resections, such that no nucleic acid purification is necessary. Appropriate procedures are described in, for example, Nuovo, G.J., 1992, "PCR *In Situ* Hybridization: Protocols And Applications", Raven Press, NY). Such analysis techniques include, DNA or RNA blotting analyses, single stranded conformational polymorphism analyses, *in situ* hybridization assays, and polymerase chain reaction analyses. Such analyses may reveal both quantitative aspects of the expression pattern of catenin p120, and qualitative aspects of catenin p120 expression and/or composition.

Alternative diagnostic methods for the detection of catenin p120 nucleic acid molecules may involve their amplification, e.g. by PCR (the experimental embodiment set forth in U.S. Patent No. 4,683,202), ligase chain reaction (Barany, 1991, Proc. Natl. Acad. Sci. USA 88:189-193), self sustained sequence replication (Guatelli *et al.*, 1990, Proc. Natl. Acad. Sci. USA 87:1874-1878), transcriptional amplification system (Kwoh *et al.*, 1989, Proc. Natl. Acad. Sci. 15 USA 86:1173-1177), Q-Beta Replicase (Lizardi *et al.*, 1988, Bio/Technology 6:1197) or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such

molecules are present in very low numbers.

Particularly suitable diagnostic methods are chip-based DNA technologies such as those described by Hacia *et al.*, 1996, Nature Genetics 14:441-447 and Shoemaker *et al.*, 1996, Nature Genetics 14:450-456. Briefly, these techniques
5 involve quantitative methods for analyzing large numbers of nucleic acid sequence targets rapidly and accurately. By tagging with oligonucleotides or using fixed probe arrays, one can employ chip technology to segregate target molecules as high density arrays and screen these molecules on the basis of hybridization.

Following detection, the results seen in a given patient may be compared with
10 a statistically significant reference group of normal patients and patients that have presenilin related pathologies. In this way, it is possible to correlate the amount or kind of catenin p120 polypeptide detected with various clinical states or predisposition to clinical states.

Substances identified according to the screening methods outlined above may
15 be formulated with standard pharmaceutically acceptable carriers and/or excipients as is routine in the pharmaceutical art. For example, a suitable substance may be dissolved in physiological saline, water or an isotonic solution for injections. The exact nature of a formulation will depend upon several factors including the particular substance to be administered and the desired route of administration. The
20 formulation will also depend on whether a pharmaceutical or veterinary use is intended. Suitable types of formulation are fully described in Remington's Pharmaceutical Sciences, Mack Publishing Company, Eastern Pennsylvania, 17th Ed. 1985, the disclosure of which is included herein of its entirety by way of reference.

The substances may be administered by enteral or parenteral routes such as
25 via oral, buccal, anal, pulmonary, intravenous, intra-arterial, intramuscular, intraperitoneal, topical, parenteral, subcutaneous, intravascular, transdermal or other appropriate administration routes.

A therapeutically effective amount of a modulator is administered to a patient. The dose of a modulator may be determined according to various
30 parameters, especially according to the substance used; the age, weight and condition of the patient to be treated; the route of administration; and the required regimen. A physician will be able to determine the required route of administration and dosage

for any particular patient. A typical daily dose is from about 0.1 to 50 mg per kg of body weight, according to the activity of the specific modulator, the age, weight and conditions of the subject to be treated, the type and severity of the degeneration and the frequency and route of administration. Preferably, daily dosage levels are from 5
5 mg to 2 g.

Modulators may have to be administered to specific sites, or otherwise targeted to brain cells. For example, the modulator may be delivered to neurons. This may be achieved, for example, by delivery via a viral strain such as herpes simplex virus. When the polynucleotide of the invention is delivered to cells by a
10 viral vector, the amount of virus administered is in the range of from 10^6 to 10^{10} pfu, preferably from 10^7 to 10^9 pfu, more preferably about 10^8 pfu for adenoviral vectors. When injected, typically 1-2ml of virus in a pharmaceutically acceptable suitable carrier or diluent is administered. The vector may comprise a promoter or other regulatory sequence that is specific to certain neurons.

15 Nucleic acid encoding catenin p120 or presenilin or a variant or fragment thereof which inhibits the presenilin/catenin p120 interaction or other catenin p120 activity may be administered to the mammal. Nucleic acid, such as RNA or DNA, and preferably, DNA, is provided in the form of a vector, such as the polynucleotides described above, which may be expressed in the cells of the mammal.

20 Nucleic acid encoding the polypeptide may be administered by any available technique. For example, the nucleic acid may be introduced by needle injection, preferably transdermally, intradermally, subcutaneously or intramuscularly. Alternatively, the nucleic acid may be delivered directly across the skin using a nucleic acid delivery device such as particle-mediated gene delivery. The nucleic
25 acid may be administered topically to the skin, or to mucosal surfaces for example by intranasal, oral, intravaginal or intrarectal administration.

Uptake of nucleic acid constructs may be enhanced by several known transfection techniques, for example those including the use of transfection agents. Examples of these agents includes cationic agents, for example, calcium phosphate
30 and DEAE-Dextran and lipofectants, for example, lipofectam and transfectam. The dosage of the nucleic acid to be administered can be altered. Typically the nucleic acid is administered in the range of 1pg to 1mg, preferably to 1pg to 10µg nucleic

acid for particle mediated gene delivery and 10 μ g to 1mg for other routes.

Where the polynucleotide giving rise to the product is under the control of an inducible promoter, it may only be necessary to induce gene expression for the duration of the treatment. Once the condition has been treated, the inducer is removed and expression of the polypeptide of the invention ceases. This will clearly have clinical advantages. Such a system may, for example, involve administering the antibiotic tetracycline, to activate gene expression via its effect on the tet repressor/VP16 fusion protein.

The use of tissue-specific promoters will be of assistance in the treatment of disease using the polypeptides, polynucleotide and vectors of the invention. It will be advantageous to be able express therapeutic genes in only the relevant affected cell types, especially where such genes are toxic when expressed in other cell types.

The routes of administration and dosages described above are intended only as a guide since a skilled physician will be able to determine readily the optimum route of administration and dosage for any particular patient and condition.

The following Examples illustrate the invention.

Example 1: Interaction of PS-1 and catenin p120 in HEK293 cells

The target protein presenilin 1 (Accession Number L42110, Hugo Number PSEN1, Unigene Number Hs.3260) was amplified by PCR using primers GGAAGTGG AAGTGG CACAGATTACCTGCACCGTTGTCC and GGAGGTTGGATTGGCTTAGATATAAAA TTGATGGAATGC to give a 1439 bp PCR product. The PCR product was cloned into an expression vector such that a single HA tag (sequence AYPYDVPDYA) was inserted at the N-terminus. This construct was transfected into HEK293 cells that were expanded under conditions selecting for the expression construct. In a representative experiment, presenilin was immunoprecipitated from 65 mg of membrane-enriched fraction derived from approximately 10⁸ transfected cells. Antibodies used were 80 μ g mouse monoclonal antibody HA11 (recognising the HA tag) (BABCO) or 30 μ g rat monoclonal antibody to presenilin 1 (Chemicon mAb 1563). Immunoprecipitates were recovered using 200 μ l of a 50/50 slurry of protein G sepharose (Pharmacia), resuspended in

sample buffer, separated on a 4-20% tris glycine gel under non-reduced conditions and stained with colloidal Coomassie blue. Bands specific to the tagged cell line were excised and in gel digested with trypsin. All trypsin digested peptides were subjected first to matrix assisted laser desorption ionization time of flight (MALDI-TOF) mass spectrometry and in a number of cases this was sufficient to make a positive identification by comparing peptide mass data with that predicted from non-redundant protein databases. In all cases peptides were further subjected to LC/MS/MS to further confirm the MALDI identification or to identify proteins not detected by MALDI-TOF, again using a non-redundant protein database. Catenin p120 was identified as interacting with presenilin 1.

Without complete sequence coverage by mass spectrometry it is not possible to inconclusively determine which p120 isoform is present. One analysis that demonstrates that the isoform present could be either 1AC or 1A (exon 20 absent, exon 18 present, presence/absence of exon 11 not established).

Example 2: Interaction of PS01 and KIAA0253 in PC12 cells, a neuronal cell line

The HA tagged human presenilin 1 construct was transfected into PC12 cells. Stable transfectants were generated and expanded under conditions selecting for the expression construct. In a representative experiment presenilin-1 was affinity-purified from 75 mg of membrane-enriched fraction derived from approximately 10^8 cells. Membrane extracts were prepared by homogenising cell pellets in a blender, taking the supernatant from a low-speed (1200 rpm/10 min) spin and subjecting it to a 100,000g/60min spin. The resulting membrane-enriched pellet was resuspended in extraction buffer and pass through a 26g needle. The solution then made 1% with dodecyl maltoside and mixed for 1 hour at 4°C. The solution was then re-spun at 100,000g for 30mins to clear precipitating debris. The antibody used was mouse monoclonal antibody HA11 (recognising the HA tag) (BABCO) covalently linked to sepharose beads (Perbio aminolink). Membrane extract was incubated with sepharose-bound antibody overnight at 4°C. Beads were then washed 6 times with extraction buffer plus 0.4mM dodecyl maltoside. Proteins that remained bound to the sepharose-antibody beads were then eluted with 100 µl sample buffer (Novex) and run on a pre-cast 4-12% 1D SDS PAGE gel in MOPS buffer (Novex) under

reducing conditions and stained with colloidal Coomassie blue. Bands specific to the tagged cell line were excised and in-gel digested with trypsin. Trypsin-digested peptides were subjected to LC/MS/MS for protein identification, using a non-redundant protein database. Catenin p120 was identified as interacting with presenilin 1 in PC12 cells.

This result extends the interaction of PS-1 and catenin p120 to a neuronal cell type that might be considered more relevant to processes involved in neurodegenerative disorders such as Alzheimer's disease.

References

- Anderton B.H. 1999. *Curr. Biol.* 9: 106-9.
- Capell A. Grunberg J. Pesold B. Diehlmann A. Citron M. Nixon R. Beyreuther K. Selkoe DJ. Haass C. 1998 *J. Biol. Chem.* 273: 3205-11.
- Daniel J.M. and Reynolds A.B. 1999. *Mol. Cell. Biol.* 19(5): 3614-23.
- Kang D.E., Soriano, S., Frosch M.P., Collins, T., Naruse, S., Sisodia, SS., Leibowitz, G., Levine, F., Koo, E.H. 1999. *J. Neurosci.* 19: 4229-37
- Keirsebilck, A., Bonneacute, S., Staes, K., Hengel, J., Nollet, F., Reynolds, A., van Roy, F. 1998. *Genomics* 50: 129-146
- Haass, C. and Mandelkow, E. 1999. *Trends in Cell Biol.* 9: 241-244
- Levesque G., Yu, G., Nishimura, M., Zhang, D.M., Levesque, L., Yu, H., Xu, D., Liang, Y., Rogaeva, E., Ikeda, M., Duthie, M., Murgolo, N., Wang, L., VanderVere, P., Bayne, M.L., Strader, C.D., Rommens, J.M., Fraser, P.E., St. George-Hyslop, P. 1999 *J. Neurochem.* 72: 999-1008
- Ray, W.J., Yao, M., Nowotny, P., Mumm., J., Zhang, W., Wu, J., Y., Kopan, R., Goate, A.M. 1999. *Proc. Natl. Acad. Sci. USA* 96: 3263-8
- Selkoe, D.J. 1998. *Trends in Cell Biol.* 8: 447-53
- Stahl, B., Diehlmann, A., Sudhof, T.C. 1999 *J. Biol. Chem.* 274: 9141-9148
- Takashima, A., Murayama, M., Murayama, O., Kohno, T., Honda, T., Yasutake, K., Nihonmatsu, N., Mercken, M., Yamaguchi, H., Sugihara, S., Wolozin, B. 1998. *Proc. Natl. Acad. Sci. USA* 9: 9637-41
- Weidemann, A., Paliga, K., Durrwang, U., Czech, C., Evin, G., Masters, C.L., Beyreuther, K. 1997. *Nat. Med.* 3: 328-32

Xia, W., Zhang, J., Perez, R., Koo, E.H., Selkoe, D.J. 1997. Proc. Natl. Acad. Sci. USA 94: 8208-13

Yu, G., Chen, F., Levesque, G., Nishimura, M., Zhang, D.M., Levesque, L., Rogaeva, E., Xu, D., Liang, Y., Duthie, M., St George-Hyslop, P.H., Fraser P.E. 1998. J. Biol. Chem. 273: 16470-5

Zhang, Z., Hartmann, H., Do, V.M., Abramowski, D., Sturchler-Pierrat, C., Staufenbiel, M., Sommer, B., van de Wetering, M., Clevers, H., Saftig, P., De Strooper, B., He, X., Yankner, B.A. 1998. Nature. 395: 698-702

Zhang, W., Han, S.W., McKeel, D.W., Goate, A., Wu, J.Y. 1998 J. Neurosci. 18: 914-22

Zhou, J., Liyanage, U., Medina, M., Ho, C., Simmons, A.D., Lovett, M., Kosik, K.S. 1997a. Neuroreport. 8: 2085-90

Zhou J. Liyanage U. Medina M. Ho C. Simmons AD. Lovett M. Kosik KS. 1997b. Neuroreport. 8: 1489-94

31
CLAIMS

1. A method of identifying a modulator of presenilin function, the method comprising:

5 (i) providing

(a) a polypeptide capable of binding a presenilin, which polypeptide comprises the amino acid sequence of SEQ ID NO: 1 or a functional variant thereof, or a fragment of either thereof which is capable of binding to presenilin;

10 (b) a presenilin or a variant thereof or a fragment of either thereof capable of binding to a polypeptide which comprises the amino acid sequence of SEQ ID NO: 1;

(c) a test substance

15 under conditions that would permit binding of a polypeptide (a) to a presenilin (b) in the absence of the test substance;

(ii) monitoring presenilin mediated activity; and

(iii) determining thereby whether the test substance is a modulator of presenilin activity.

20 2. A method according to claim 1 wherein step (ii) comprises monitoring the interaction between polypeptide (a) and presenilin (b).

3. A method according to claim 2 wherein the modulator inhibits the binding of the polypeptide (a) to the presenilin (b).

4. A method according to claim 2 wherein the modulator enhances the binding of the polypeptide (a) to the presenilin (b).

25 5. A method according to any one of the preceding claims wherein the presenilin (b) is presenilin 1 or a said fragment thereof.

6. A method according to any one of the preceding claims wherein step (i) comprises:

30 (a) transfecting a cell with polynucleotides encoding a polypeptide (a) and a presenilin(b);

(b) allowing the said cell to express the polypeptide (a) and presenilin (b); and

- (c) contacting the said cell with a test substance.

7. A method according to any one of claims 1 to 5 wherein step (i) comprises:

- (a) transfecting a cell with polynucleotides encoding a polypeptide(a), a presenilin (b) and a test substance (c) which is a peptide and
- (b) allowing the said cell to express the polypeptide (a), presenilin (b) and the peptide test substance (c).

8. A method according to any one of claims 1 to 5 wherein step (i) comprises:

- (a) transfecting a first cell with a polynucleotide encoding a polypeptide (a);
- (b) transfecting a second cell with a polynucleotide encoding a presenilin (b);
- (c) allowing said first cell to express the polypeptide(a) and said second cell to express the presenilin (b);
- (d) preparing a cell extract from each of said first and second cell; and
- (e) contacting said cell extract from first cell and said cell extract from second cell in the presence of a test substance.

9. A method for identification of a compound that modulates catenin p120 activity, which method comprises:

- (i) contacting a catenin p120 polypeptide comprising
- (a) the amino acid sequence of SEQ ID NO: 1; or
- (b) a variant thereof or a fragment of either thereof which maintains a catenin p120 function; with a test compound and
- (ii) monitoring for catenin p120 activity thereby determining whether the test compound is a modulator of catenin p120.

10. A method according to claim 9 wherein the catenin p120 activity comprises the ability of the polypeptide to interact with a presenilin, or a variant thereof or a fragment of either thereof.

11. A modulator identified by a method according to any one of the preceding claims for use in a method of treatment of the human or animal body by therapy.

12. Use of a modulator identified by a method according to any one of claims 1 to 10 in the manufacture of a medicament for the treatment or prophylaxis of a disorder that is responsive to modulation of presenilin activity.

13. Use of a polypeptide capable of binding a presenilin, which polypeptide comprises the amino acid sequence of SEQ ID NO: 1 or a variant or fragment of SEQ ID NO: 1 which is capable of binding a presenilin in the manufacture of a medicament for the treatment, prophylaxis or diagnosis of a disorder that is responsive to modulation of presenilin activity.

14. Use of a polynucleotide which encodes a polypeptide as defined in claim 13 or a catenin p120 polypeptide as defined in claim 9 comprising:

- (a) the sequence of SEQ ID NO: 2; or
- (b) a sequence that hybridizes to the complement of SEQ ID NO: 2; or
- (c) a sequence that is degenerate as a result of the genetic code with respect to a sequence defined in (a) or (b); or
- (d) a sequence that is complementary to a polynucleotide defined in (a), (b) or (c);

in the manufacture of a medicament for the treatment, prophylaxis or diagnosis of a disorder that is responsive to modulation of presenilin activity.

15. Use according to any one of claims 12 to 14 wherein the disorder is Alzheimer's disease.

16. A method of treatment, prophylaxis or diagnosis of Alzheimer's disease which method comprises administering an effective amount of a polypeptide as defined in claim 13, a polynucleotide as defined in claim 14 or a modulator identified by a method according to any one of claims 1 to 10 to a human or animal in need of such treatment.

SEQUENCE LISTING

<110> GLAXO GROUP LIMITED

<120> ASSAY

<130> N79188A GCW/SER

<140>

<141>

<150> GB 0005895.8

<151> 2000-03-10

<160> 6

<170> Patent In Ver. 2.1

<210> 1

<211> 939

<212> PRT

<213> Homo sapiens

<400> 1

```

Met Asp Asp Ser Glu Val Glu Ser Thr Ala Ser Ile Leu Ala Ser Val
 1           5           10           15
Lys Glu Gln Glu Ala Gln Phe Glu Lys Leu Thr Arg Ala Leu Glu Glu
      20           25           30
Glu Arg Arg His Val Ser Ala Gln Leu Glu Arg Val Arg Val Ser Pro
      35           40           45
Gln Asp Ala Asn Pro Leu Met Ala Asn Gly Thr Leu Thr Arg Arg His
      50           55           60
Gln Asn Gly Arg Phe Val Gly Asp Ala Asp Leu Glu Arg Gln Lys Phe
      65           70           75           80
Ser Asp Leu Lys Leu Asn Gly Pro Gln Asp His Ser His Leu Leu Tyr
      85           90           95
Ser Thr Ile Pro Arg Met Gln Glu Pro Gly Gln Ile Val Glu Thr Tyr
      100          105          110
Thr Glu Glu Asp Pro Glu Gly Ala Met Ser Val Val Ser Val Glu Thr
      115          120          125
Ser Asp Asp Gly Thr Thr Arg Arg Thr Glu Thr Thr Val Lys Lys Val
      130          135          140
Val Lys Thr Val Thr Thr Arg Thr Val Gln Pro Val Ala Met Gly Pro
      145          150          155          160
Asp Gly Leu Pro Val Asp Ala Ser Ser Val Ser Asn Asn Tyr Ile Gln
      165          170          175
Thr Leu Gly Arg Asp Phe Arg Lys Asn Gly Asn Gly Gly Pro Gly Pro
      180          185          190
Tyr Val Gly Gln Ala Gly Thr Ala Thr Leu Pro Arg Asn Phe His Tyr
      195          200          205
Pro Pro Asp Gly Tyr Ser Arg His Tyr Glu Asp Gly Tyr Pro Gly Gly
      210          215          220
Ser Asp Asn Tyr Gly Ser Leu Ser Arg Val Thr Arg Ile Glu Glu Arg
      225          230          235          240
Tyr Arg Pro Ser Met Glu Gly Tyr Arg Ala Pro Ser Arg Gln Asp Val
      245          250          255

Tyr Gly Pro Gln Pro Gln Val Arg Val Gly Gly Ser Ser Val Asp Leu

```

260 265 270
 His Arg Phe His Pro Glu Pro Tyr Gly Leu Glu Asp Asp Gln Arg Ser
 275 280 285
 Met Gly Tyr Asp Asp Leu Asp Tyr Gly Met Met Ser Asp Tyr Gly Thr
 290 295 300
 Ala Arg Arg Thr Gly Thr Pro Ser Asp Pro Arg Arg Arg Leu Arg Ser
 305 310 315 320
 Tyr Glu Asp Met Ile Gly Glu Glu Val Pro Ser Asp Gln Tyr Tyr Trp
 325 330 335
 Ala Pro Leu Ala Gln His Glu Arg Gly Ser Leu Ala Ser Leu Asp Ser
 340 345 350
 Leu Arg Lys Gly Gly Pro Pro Pro Pro Asn Trp Arg Gln Pro Glu Leu
 355 360 365
 Pro Glu Val Ile Ala Met Leu Gly Phe Arg Leu Asp Ala Val Lys Ser
 370 375 380
 Asn Ala Ala Ala Tyr Leu Gln His Leu Cys Tyr Arg Asn Asp Lys Val
 385 390 395 400
 Lys Thr Asp Val Arg Lys Leu Lys Gly Ile Pro Val Leu Val Gly Leu
 405 410 415
 Leu Asp His Pro Lys Lys Glu Val His Leu Gly Ala Cys Gly Ala Leu
 420 425 430
 Lys Asn Ile Ser Phe Gly Arg Asp Gln Asp Asn Lys Ile Ala Ile Lys
 435 440 445
 Asn Cys Asp Gly Val Pro Ala Leu Val Arg Leu Leu Arg Lys Ala Arg
 450 455 460
 Asp Met Asp Leu Thr Glu Val Ile Thr Gly Thr Leu Trp Asn Leu Ser
 465 470 475 480
 Ser His Asp Ser Ile Lys Met Glu Ile Val Asp His Ala Leu His Ala
 485 490 495
 Leu Thr Asp Glu Val Ile Ile Pro His Ser Gly Trp Glu Arg Glu Pro
 500 505 510
 Asn Glu Asp Cys Lys Pro Arg His Ile Glu Trp Glu Ser Val Leu Thr
 515 520 525
 Asn Thr Ala Gly Cys Leu Arg Asn Val Ser Ser Glu Arg Ser Glu Ala
 530 535 540
 Arg Arg Lys Leu Arg Glu Cys Asp Gly Leu Val Asp Ala Leu Ile Phe
 545 550 555 560
 Ile Val Gln Ala Glu Ile Gly Gln Lys Asp Ser Asp Ser Lys Leu Val
 565 570 575
 Glu Asn Cys Val Cys Leu Leu Arg Asn Leu Ser Tyr Gln Val His Arg
 580 585 590
 Glu Ile Pro Gln Ala Glu Arg Tyr Gln Glu Ala Ala Pro Asn Val Ala
 595 600 605
 Asn Asn Thr Gly Pro His Ala Ala Ser Cys Phe Gly Ala Lys Lys Gly
 610 615 620
 Lys Asp Glu Trp Phe Ser Arg Gly Lys Lys Pro Ile Glu Asp Pro Ala
 625 630 635 640
 Asn Asp Thr Val Asp Phe Pro Lys Arg Thr Ser Pro Ala Arg Gly Tyr
 645 650 655
 Glu Leu Leu Phe Gln Pro Glu Val Val Arg Ile Tyr Ile Ser Leu Leu
 660 665 670
 Lys Glu Ser Lys Thr Pro Ala Ile Leu Glu Ala Ser Ala Gly Ala Ile
 675 680 685

 Gln Asn Leu Cys Ala Gly Arg Trp Thr Tyr Gly Arg Tyr Ile Arg Ser
 690 695 700
 Ala Leu Arg Gln Glu Lys Ala Leu Ser Ala Ile Ala Asp Leu Leu Thr

cgg cgc cac gtc tcg gcg cag ctg gaa cgc gtc cgg gtc tca cca caa 583
 Arg Arg His Val Ser Ala Gln Leu Glu Arg Val Arg Val Ser Pro Gln
 35 40 45
 gat gcc aac cca ctc atg gcc aac ggc aca ctc acc cgc cgg cat cag 731
 Asp Ala Asn Pro Leu Met Ala Asn Gly Thr Leu Thr Arg Arg His Gln
 50 55 60 65
 aac ggc cgg ttt gtg ggc gat gct gac ctt gaa aga cag aaa ttt tca 779
 Asn Gly Arg Phe Val Gly Asp Ala Asp Leu Glu Arg Gln Lys Phe Ser
 70 75 80
 gat ttg aaa ctc aac gga ccc cag gat cac agt cac ctt cta tat agc 827
 Asp Leu Lys Leu Asn Gly Pro Gln Asp His Ser His Leu Leu Tyr Ser
 85 90 95
 acc atc ccc agg atg cag gag ccg ggg cag att gtg gag acc tac acg 875
 Thr Ile Pro Arg Met Gln Glu Pro Gly Gln Ile Val Glu Thr Tyr Thr
 100 105 110
 gag gag gat cct gag gga gcc atg tct gta gtc tct gtg gag acc tca 923
 Glu Glu Asp Pro Glu Gly Ala Met Ser Val Val Ser Val Glu Thr Ser
 115 120 125
 gat gat ggg acc act cgg cgc aca gag acc acg gtc aag aaa gta gtg 971
 Asp Asp Gly Thr Thr Arg Arg Thr Glu Thr Thr Val Lys Lys Val Val
 130 135 140 145
 aag act gtg aca aca cgg aca gta cag cca gtc gct atg gga cca gac 1019
 Lys Thr Val Thr Thr Arg Thr Val Gln Pro Val Ala Met Gly Pro Asp
 150 155 160
 ggg ttg cct gtg gat gct tca tca gtt tct aac aac tat atc cag act 1067
 Gly Leu Pro Val Asp Ala Ser Ser Val Ser Asn Asn Tyr Ile Gln Thr
 165 170 175
 ttg ggt cgt gat ttc cgc aag aat ggc aat ggg gga cct ggt ccc tat 1115
 Leu Gly Arg Asp Phe Arg Lys Asn Gly Asn Gly Gly Pro Gly Pro Tyr
 180 185 190
 gtg ggg caa gct ggc act gct acc ctt cct agg aac ttc cac tac cct 1163
 Val Gly Gln Ala Gly Thr Ala Thr Leu Pro Arg Asn Phe His Tyr Pro
 195 200 205
 cct gat ggt tat agt cgc cac tat gaa gat ggt tat cca ggt ggc agt 1211
 Pro Asp Gly Tyr Ser Arg His Tyr Glu Asp Gly Tyr Pro Gly Gly Ser
 210 215 220 225
 gat aac tat ggc agt ctg tcc cgg gtg acc cgc att gag gag cgg tat 1259
 Asp Asn Tyr Gly Ser Leu Ser Arg Val Thr Arg Ile Glu Glu Arg Tyr
 230 235 240
 agg ccc agc atg gaa ggc tac cgg gca cct agt aga cag gat gtg tat 1307
 Arg Pro Ser Met Glu Gly Tyr Arg Ala Pro Ser Arg Gln Asp Val Tyr
 245 250 255
 ggg ccc caa ccc cag gtt cgg gta ggt ggg agc agc gtg gat ctg cat 1355
 Gly Pro Gln Pro Gln Val Arg Val Gly Gly Ser Ser Val Asp Leu His
 260 265 270
 cgc ttt cat cca gag cct tat ggg cta gag gat gac cag cgt agt atg 1403
 Arg Phe His Pro Glu Pro Tyr Gly Leu Glu Asp Asp Gln Arg Ser Met
 275 280 285
 ggc tat gat gac ctg gat tat ggt atg atg tct gat tat ggc act gcc 1451
 Gly Tyr Asp Asp Leu Asp Tyr Gly Met Met Ser Asp Tyr Gly Thr Ala
 290 295 300 305
 cgt cgg act ggg aca ccc tct gac cct cgt cgg cgc ctc agg agc tat 1499
 Arg Arg Thr Gly Thr Pro Ser Asp Pro Arg Arg Arg Leu Arg Ser Tyr
 310 315 320
 gaa gac atg att ggt gag gag gtg cca tcg gat caa tac tac tgg gct 1547
 Glu Asp Met Ile Gly Glu Glu Val Pro Ser Asp Gln Tyr Tyr Trp Ala
 325 330 335
 cct ttg gcc cag cat gag cga gga agt tta gca agc ttg gat agc ctg 1595

Pro Leu Ala Gln His Glu Arg Gly Ser Leu Ala Ser Leu Asp Ser Leu
 340 345 350
 cgc aaa gga ggg cct cca cct aat tgg aga cag cca gag ctg cca 1643
 Arg Lys Gly Gly Pro Pro Pro Pro Asn Trp Arg Gln Pro Glu Leu Pro
 355 360 365
 gag gtg atc gcc atg ctt gga ttc cgc ttg gat gct gtc aag tcc aat 1691
 Glu Val Ile Ala Met Leu Gly Phe Arg Leu Asp Ala Val Lys Ser Asn
 370 375 380 385
 gca gct gca tac ctg caa cac tta tgc tac cgc aat gac aag gtg aag 1739
 Ala Ala Ala Tyr Leu Gln His Leu Cys Tyr Arg Asn Asp Lys Val Lys
 390 395 400
 act gac gtg cgg aag ctc aag ggc atc cca gta ctg gtg gga ttg tta 1787
 Thr Asp Val Arg Lys Leu Lys Gly Ile Pro Val Leu Val Gly Leu Leu
 405 410 415
 gac cat ccc aaa aag gaa gtg cac ctt gga gcc tgt gga gct ctc aag 1835
 Asp His Pro Lys Lys Glu Val His Leu Gly Ala Cys Gly Ala Leu Lys
 420 425 430
 aat atc tct ttt gga cgt gac cag gat aac aag att gcc ata aaa aac 1883
 Asn Ile Ser Phe Gly Arg Asp Gln Asp Asn Lys Ile Ala Ile Lys Asn
 435 440 445
 tgt gat ggt gtg cct gcc ctt gtg cga ttg ctt cga aag gct cgt gat 1931
 Cys Asp Gly Val Pro Ala Leu Val Arg Leu Leu Arg Lys Ala Arg Asp
 450 455 460 465
 atg gac ctt act gaa gtt att acc gga acc ctg tgg aat ctt tca tcc 1979
 Met Asp Leu Thr Glu Val Ile Thr Gly Thr Leu Trp Asn Leu Ser Ser
 470 475 480
 cat gac tca atc aaa atg gag att gtg gac cat gca ctg cat gcc ttg 2027
 His Asp Ser Ile Lys Met Glu Ile Val Asp His Ala Leu His Ala Leu
 485 490 495
 aca gat gaa gtg atc att cct cat tct ggt tgg gag cgg gaa cct aat 2075
 Thr Asp Glu Val Ile Ile Pro His Ser Gly Trp Glu Arg Glu Pro Asn
 500 505 510
 gaa gac tgt aag cca cgc cat att gag tgg gaa tcg gtg ctc acc aac 2123
 Glu Asp Cys Lys Pro Arg His Ile Glu Trp Glu Ser Val Leu Thr Asn
 515 520 525
 aca gct ggc tgc ctt agg aat gta agc tca gag agg agt gaa gct cgc 2171
 Thr Ala Gly Cys Leu Arg Asn Val Ser Ser Glu Arg Ser Glu Ala Arg
 530 535 540 545
 cgg aaa ctt cgg gaa tgt gat ggt tta gtt gat gcc ctc att ttc att 2219
 Arg Lys Leu Arg Glu Cys Asp Gly Leu Val Asp Ala Leu Ile Phe Ile
 550 555 560
 gtt cag gct gag att ggg cag aag gat tca gac agc aag ctt gta gag 2267
 Val Gln Ala Glu Ile Gly Gln Lys Asp Ser Asp Ser Lys Leu Val Glu
 565 570 575
 aac tgt gtt tgc ctt ctt cgg aac tta tca tat caa gtt cac cgg gag 2315
 Asn Cys Val Cys Leu Leu Arg Asn Leu Ser Tyr Gln Val His Arg Glu
 580 585 590
 atc cca cag gca gag cgt tac caa gag gca gct ccc aat gtt gcc aac 2363
 Ile Pro Gln Ala Glu Arg Tyr Gln Glu Ala Ala Pro Asn Val Ala Asn
 595 600 605
 aat act ggg cca cat gct gcc agt tgc ttt ggg gcc aag aag ggc aaa 2411
 Asn Thr Gly Pro His Ala Ala Ser Cys Phe Gly Ala Lys Lys Gly Lys
 610 615 620 625
 gat gag tgg ttc tcc aga ggg aaa aaa cct ata gag gat cca gca aac 2459
 Asp Glu Trp Phe Ser Arg Gly Lys Lys Pro Ile Glu Asp Pro Ala Asn
 630 635 640
 gat aca gtg gat ttc cct aaa aga acg agt cca gct cga ggc tat gag 2507
 Asp Thr Val Asp Phe Pro Lys Arg Thr Ser Pro Ala Arg Gly Tyr Glu

```

        645              650              655
ctc tta ttt cag cca gag gtg gtt cgg ata tac atc tca ctt ctt aag 2555
Leu Leu Phe Gln Pro Glu Val Val Arg Ile Tyr Ile Ser Leu Leu Lys

        660              665              670
gag agc aag act cct gcc atc cta gaa gcc tca gct gga gct atc cag 2603
Glu Ser Lys Thr Pro Ala Ile Leu Glu Ala Ser Ala Gly Ala Ile Gln

        675              680              685
aac ttg tgt gct ggg cgc tgg acg tat ggt cga tac atc cgc tct gct 2651
Asn Leu Cys Ala Gly Arg Trp Thr Tyr Gly Arg Tyr Ile Arg Ser Ala

        690              695              700              705
ctg cgt caa gag aag gct ctt tct gcc ata gct gac ctc ctg act aat 2699
Leu Arg Gln Glu Lys Ala Leu Ser Ala Ile Ala Asp Leu Leu Thr Asn

        710              715              720
gaa cat gaa cgg gtg gtg aaa gct gca tct gga gca ctg aga aac ctg 2747
Glu His Glu Arg Val Val Lys Ala Ala Ser Gly Ala Leu Arg Asn Leu

        725              730              735
gct gtg gat gct cgc aac aaa gaa tta att ggt aaa cat gct att cct 2795
Ala Val Asp Ala Arg Asn Lys Glu Leu Ile Gly Lys His Ala Ile Pro

        740              745              750
aac ttg gta aag aat ctg cca gga gga cag cag aac tcc tct tgg aat 2843
Asn Leu Val Lys Asn Leu Pro Gly Gly Gln Gln Asn Ser Ser Trp Asn

        755              760              765
ttc tct gag gac act gtc atc tct att ttg aac act atc aac gag gtt 2891
Phe Ser Glu Asp Thr Val Ile Ser Ile Leu Asn Thr Ile Asn Glu Val

        770              775              780              785
atc gct gag aac ttg gag gct gcc aaa aag ctt cga gag aca cag ggt 2939
Ile Ala Glu Asn Leu Glu Ala Ala Lys Lys Leu Arg Glu Thr Gln Gly

        790              795              800
att gag aag ctg gtg ttg atc aac aaa tca ggg aac cgc tca gaa aaa 2987
Ile Glu Lys Leu Val Leu Ile Asn Lys Ser Gly Asn Arg Ser Glu Lys

        805              810              815
gaa gtt cga gca gca gca ctt gta tta cag aca atc tgg gga tat aag 3035
Glu Val Arg Ala Ala Ala Leu Val Leu Gln Thr Ile Trp Gly Tyr Lys

        820              825              830
gaa ctg cgg aag cca ctg gaa aaa gaa gga tgg aag aaa tca gac ttt 3083
Glu Leu Arg Lys Pro Leu Glu Lys Glu Gly Trp Lys Lys Ser Asp Phe

        835              840              845
cag gtg aat cta aac aat gct tcc cga agc cag agc agt cat tca tat 3131
Gln Val Asn Leu Asn Asn Ala Ser Arg Ser Gln Ser Ser His Ser Tyr

        850              855              860              865
gat gat agt act ctc cct ctc att gac cgg aac caa aaa tca gat aag 3179
Asp Asp Ser Thr Leu Pro Leu Ile Asp Arg Asn Gln Lys Ser Asp Lys

        870              875              880
aaa cct gat cgg gaa gaa att cag atg agc aat atg gga tca aac aca 3227
Lys Pro Asp Arg Glu Glu Ile Gln Met Ser Asn Met Gly Ser Asn Thr

        885              890              895
aaa tca cta gat aac aac tat tcc aca cca aat gag aga gga gac cac 3275
Lys Ser Leu Asp Asn Asn Tyr Ser Thr Pro Asn Glu Arg Gly Asp His

        900              905              910
aat aga aca ctg gat cga tcg ggg gat cta ggc gac atg gag cca ttg 3323
Asn Arg Thr Leu Asp Arg Ser Gly Asp Leu Gly Asp Met Glu Pro Leu

        915              920              925
aag gga aca aca ccc ttg atg cag aag att tag caccactatc tccgttccat 3376
Lys Gly Thr Thr Pro Leu Met Gln Lys Ile

        930              935              940
ctgggcttat atgtactttt attttttggt ggtgaaattg actgatgatt ttcctttttc 3436
ttcgtggac tattgtgcca actgccaggc tgctctctgc ccttacagcc ctaagtggct 3496
gccttctttc catcaactcc caacttcttc ctgtgaagtt taattgtctc aacgcctccc 3556

```


cctccccat tccctccatt tttctcccga gaaacctgac tcaattatgt gcatatattg 3616
 agaaactgct gcagattagt tctttttgcc agttttccct ggaactcctg gccttttgtg 3676
 gaggggaggg atggagagaa taggaatctt cactagaagc cgtgggaaga attggaagtt 3736
 acatgctgta tatgcaatgt ccagcagctt gataaactga cgattcttaa tcaagatttt 3796
 tttcctgatg ggggaaggac ttttattttc ttttagagag gggaaaagtgt gagctcttcc 3856
 cttattccta atggctatgt ttgaagcaaa gaaggccagc aacattggca catgccacct 3916
 ggcaaggac ccttgagtaa gtgaaggtct cctaaaactg ggattaagaa accttgctct 3976
 cctcatctcc aaggcagggg ccatcaagaa cctacagact ccatctcttc tgcaagcctc 4036
 atgccaacct rgggtatgt ctgctgcccc ttaaaccacag gctgtcctta acccactctt 4096
 cctgcccctg gatatgtctg ctgagttggc ctggccattt ccaagaggct gtagaaaggg 4156
 gagaatgca aggaagactt ttggtagaga aggagcagaa agatgtgttt ttgggaagaa 4216
 gaagacctct agggagagct agtaggaatg tacatgaagc aattagtctg aaactggctt 4276
 cccactctcc ccgtttctcc ttttctatc cttataggcc tgtcccttgc ctctgcccctg 4336
 gattggttgg caaactatag gacttgatgt acataactcc tgtccctttt cccttacaag 4396
 gtggggattg cccctggctt tgcctctctt ttgtgccttt ggccctgggt gcattctctc 4456
 ccgcccctcc atgtgccttt ctttgcctct gcagctctcat ttctcataat ttgcaaat 4516
 atattttgtt gctttcttac ctactattgg ccctaaatag cagaaagaag agaagtgacc 4576
 gagagaacct cagattcttc attgaggatt ggtatagcca tgatttcagt catagcaagc 4636
 ttttgcctaa cagcatatgg gtgggatttg gcaaaaatcc tattctgatg aatctcaaag 4696
 taaggctggt aagagaagtg agtgggtgta ctcttactcc ttaggtgccc agaatttacc 4756
 atcatctctg aaggagttac aggggaagtgg tctccccaat tctccctccc ctccagttatt 4816
 gccccccttc acttttagcat atattaatta gcagggtggg ctagagaat cagctgctat 4876
 gcgggtgat tattattatt atttctaata cttttcctta ttgtccttct actcccttca 4936
 atctaatacta aaagctctgt tccatgcaac tggagttcct tatccctctc tccccctccc 4996
 cttatatatt gaggtatgg ggtaggagaa aagtgcacaa cccaccaccc cctctactcg 5056
 tgcattaaaa tttcttattt acccttttcc cccttcccat ttcttccac ttctatctac 5116
 cttttctggc aaaaaggagc cttttgctct ctgtgacctt aagagcacac tgcacaggga 5176
 aaattgcccc atccagacct ggctccactc ttgatctctc ttgtcctctt ctgctctttt 5236
 cctggtgctc ttttttctcg gtgggtgtg ggtaatagaa cagccgtggg cttttgggga 5296
 cctttaactt ttttttctct cttttgttta taaaaaacac taaacattca attccagaga 5356
 acccaaaatc ccacttctcc accgaacact actaaggggc ttgtgttctg ctccatactt 5416
 tttctctttt ctttctgtct tgttaatgct tttaaaaaca aatgagtttt ttatataaat 5476
 aaagttttta aagtgtgtat gtgggggggc tgtgtcattt cttcacttca agctgttatt 5536
 ttttccctgc tttgcatctt tgttacttcc ttatgtatca gtgtccttcc cagagcaacc 5596
 agaaggaggt tataccagga tttattttga gctcagcccc aactctttat caagcaacat 5656
 tcttggttaac tatatgtgaa acattttttc tttgaagat ttttaaaaat tgaatgtggc 5716
 tgaagttgaa catgggagct tattgctaatt tttagagatag gaaactgaag cataaagaat 5776
 taatgactta ctttaattac tggaaattctt ctgcaacatt tgacaaaact aaccttgaat 5836
 aaggccact gtaatacgtg gctctcttaa atataacact taggactaga agattagaaa 5896
 ctaccaatcc caactacgta ataggaaaat gtaggatcaa aaggcccatg tatataagta 5956
 ctgaccactg ggcataatg ttgcttctca ggctatatgc agtcctttag tcagaagtca 6016
 ataggcctat ttattaatat ttacagacc atattacctg gattaccagg gactatcttt 6076
 gctgcagaga tcaagggtta agatctatgg gaagatactt atttttctga ggctcttatg 6136
 tcctgtcata taattaaaga ctcaagagaa tttatgtgaa atgctttctg tatgccccaa 6196
 tcttttagatt aaaattatat acctgctcct 6226

<210> 3

<211> 467

<212> PRT

<213> Homo sapiens

<400> 3

Met Thr Glu Leu Pro Ala Pro Leu Ser Tyr Phe Gln Asn Ala Gln Met
 1 5 10 15
 Ser Glu Asp Asn His Leu Ser Asn Thr Val Arg Ser Gln Asn Asp Asn
 20 25 30
 Arg Glu Arg Gln Glu His Asn Asp Arg Arg Ser Leu Gly His Pro Glu

35 40 45
 Pro Leu Ser Asn Gly Arg Pro Gln Gly Asn Ser Arg Gln Val Val Glu
 50 55 60
 Gln Asp Glu Glu Glu Asp Glu Glu Leu Thr Leu Lys Tyr Gly Ala Lys
 65 70 75 80
 His Val Ile Met Leu Phe Val Pro Val Thr Leu Cys Met Val Val Val
 85 90 95
 Val Ala Thr Ile Lys Ser Val Ser Phe Tyr Thr Arg Lys Asp Gly Gln
 100 105 110
 Leu Ile Tyr Thr Pro Phe Thr Glu Asp Thr Glu Thr Val Gly Gln Arg
 115 120 125
 Ala Leu His Ser Ile Leu Asn Ala Ala Ile Met Ile Ser Val Ile Val
 130 135 140
 Val Met Thr Ile Leu Leu Val Val Leu Tyr Lys Tyr Arg Cys Tyr Lys
 145 150 155 160
 Val Ile His Ala Trp Leu Ile Ile Ser Ser Leu Leu Leu Phe Phe
 165 170 175
 Phe Ser Phe Ile Tyr Leu Gly Glu Val Phe Lys Thr Tyr Asn Val Ala
 180 185 190
 Val Asp Tyr Ile Thr Val Ala Leu Leu Ile Trp Asn Phe Gly Val Val
 195 200 205
 Gly Met Ile Ser Ile His Trp Lys Gly Pro Leu Arg Leu Gln Gln Ala
 210 215 220
 Tyr Leu Ile Met Ile Ser Ala Leu Met Ala Leu Val Phe Ile Lys Tyr
 225 230 235 240
 Leu Pro Glu Trp Thr Ala Trp Leu Ile Leu Ala Val Ile Ser Val Tyr
 245 250 255
 Asp Leu Val Ala Val Leu Cys Pro Lys Gly Pro Leu Arg Met Leu Val
 260 265 270
 Glu Thr Ala Gln Glu Arg Asn Glu Thr Leu Phe Pro Ala Leu Ile Tyr
 275 280 285
 Ser Ser Thr Met Val Trp Leu Val Asn Met Ala Glu Gly Asp Pro Glu
 290 295 300
 Ala Gln Arg Arg Val Ser Lys Asn Ser Lys Tyr Asn Ala Glu Ser Thr
 305 310 315 320
 Glu Arg Glu Ser Gln Asp Thr Val Ala Glu Asn Asp Asp Gly Gly Phe
 325 330 335
 Ser Glu Glu Trp Glu Ala Gln Arg Asp Ser His Leu Gly Pro His Arg
 340 345 350
 Ser Thr Pro Glu Ser Arg Ala Ala Val Gln Glu Leu Ser Ser Ser Ile
 355 360 365
 Leu Ala Gly Glu Asp Pro Glu Glu Arg Gly Val Lys Leu Gly Leu Gly
 370 375 380
 Asp Phe Ile Phe Tyr Ser Val Leu Val Gly Lys Ala Ser Ala Thr Ala
 385 390 395 400
 Ser Gly Asp Trp Asn Thr Thr Ile Ala Cys Phe Val Ala Ile Leu Ile
 405 410 415
 Gly Leu Cys Leu Thr Leu Leu Leu Leu Ala Ile Phe Lys Lys Ala Leu
 420 425 430
 Pro Ala Leu Pro Ile Ser Ile Thr Phe Gly Leu Val Phe Tyr Phe Ala
 435 440 445
 Thr Asp Tyr Leu Val Gln Pro Phe Met Asp Gln Leu Ala Phe His Gln
 450 455 460
 Phe Tyr Ile
 465

<210> 4

<211> 2765

<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (249)..(1652)

<400> 4

```

tgggacaggc agctccgggg tccgcggttt cacatcggaa aaaaaacagc ggctgggtctg 60
gaaggaaacct gagctacgag ccgcggcggc agcggggcgg cggggaagcg tatacctaata 120
ctgggagacct gcaagtgaca acagcctttg cggtccttag acagcttggc ctggaggaga 180
acacatgaaa gaaagaacct caagaggctt tgttttctgt gaaacagtat ttctatacag 240

ttgctcca atg aca gag tta cct gca ccg ttg tcc tac ttc cag aat gca 290
      Met Thr Glu Leu Pro Ala Pro Leu Ser Tyr Phe Gln Asn Ala
      1          5          10
cag atg tct gag gac aac cac ctg agc aat act gta cgt agc cag aat 338
Gln Met Ser Glu Asp Asn His Leu Ser Asn Thr Val Arg Ser Gln Asn
      15          20          25          30
gac aat aga gaa cgg cag gag cac aac gac aga cgg agc ctt ggc cac 386
Asp Asn Arg Glu Arg Gln Glu His Asn Asp Arg Arg Ser Leu Gly His
      35          40          45
cct gag cca tta tct aat gga cga ccc cag ggt aac tcc cgg cag gtg 434
Pro Glu Pro Leu Ser Asn Gly Arg Pro Gln Gly Asn Ser Arg Gln Val
      50          55          60
gtg gag caa gat gag gaa gaa gat gag gag ctg aca ttg aaa tat ggc 482
Val Glu Gln Asp Glu Glu Glu Asp Glu Glu Leu Thr Leu Lys Tyr Gly
      65          70          75
gcc aag cat gtg atc atg ctc ttt gtc cct gtg act ctc tgc atg gtg 530
Ala Lys His Val Ile Met Leu Phe Val Pro Val Thr Leu Cys Met Val
      80          85          90
gtg gtc gtg gct acc att aag tca gtc agc ttt tat acc cgg aag gat 578
Val Val Val Ala Thr Ile Lys Ser Val Ser Phe Tyr Thr Arg Lys Asp
      95          100          105          110
ggg cag cta atc tat acc cca ttc aca gaa gat acc gag act gtg ggc 626
Gly Gln Leu Ile Tyr Thr Pro Phe Thr Glu Asp Thr Glu Thr Val Gly
      115          120          125
cag aga gcc ctg cac tca att ctg aat gct gcc atc atg atc agt gtc 674
Gln Arg Ala Leu His Ser Ile Leu Asn Ala Ala Ile Met Ile Ser Val
      130          135          140
att gtt gtc atg act atc ctc ctg gtt ctg tat aaa tac agg tgc 722
Ile Val Val Met Thr Ile Leu Leu Val Val Leu Tyr Lys Tyr Arg Cys
      145          150          155
tat aag gtc atc cat gcc tgg ctt att ata tca tct cta ttg ttg ctg 770
Tyr Lys Val Ile His Ala Trp Leu Ile Ile Ser Ser Leu Leu Leu Leu
      160          165          170
ttc ttt ttt tca ttc att tac ttg ggg gaa gtg ttt aaa acc tat aac 818
Phe Phe Phe Ser Phe Ile Tyr Leu Gly Glu Val Phe Lys Thr Tyr Asn
      175          180          185          190
gtt gct gtg gac tac att act gtt gca ctc ctg atc tgg aat ttt ggt 866
Val Ala Val Asp Tyr Ile Thr Val Ala Leu Leu Ile Trp Asn Phe Gly
      195          200          205
gtg gtg gga atg att tcc att cac tgg aaa ggt cca ctt cga ctc cag 914
Val Val Gly Met Ile Ser Ile His Trp Lys Gly Pro Leu Arg Leu Gln
      210          215          220
cag gca tat ctc att atg att agt gcc ctc atg gcc ctg gtg ttt atc 962
Gln Ala Tyr Leu Ile Met Ile Ser Ala Leu Met Ala Leu Val Phe Ile
      225          230          235

```

aag tac ctc cct gaa tgg act gcg tgg ctc atc ttg gct gtg att tca 1010
 Lys Tyr Leu Pro Glu Trp Thr Ala Trp Leu Ile Leu Ala Val Ile Ser
 240 245 250
 gta tat gat tta gtg gct gtt ttg tgt ccg aaa ggt cca ctt cgt atg 1058
 Val Tyr Asp Leu Val Ala Val Leu Cys Pro Lys Gly Pro Leu Arg Met
 255 260 265 270
 ctg gtt gaa aca gct cag gag aga aat gaa acg ctt ttt cca gct ctc 1106
 Leu Val Glu Thr Ala Gln Glu Arg Asn Glu Thr Leu Phe Pro Ala Leu
 275 280 285
 att tac tcc tca aca atg gtg tgg ttg gtg aat atg gca gaa gga gac 1154
 Ile Tyr Ser Ser Thr Met Val Trp Leu Val Asn Met Ala Glu Gly Asp
 290 295 300
 ccg gaa gct caa agg aga gta tcc aaa aat tcc aag tat aat gca gaa 1202
 Pro Glu Ala Gln Arg Arg Val Ser Lys Asn Ser Lys Tyr Asn Ala Glu
 305 310 315
 agc aca gaa agg gag tca caa gac act gtt gca gag aat gat gat ggc 1250
 Ser Thr Glu Arg Glu Ser Gln Asp Thr Val Ala Glu Asn Asp Asp Gly
 320 325 330
 ggg ttc agt gag gaa tgg gaa gcc cag agg gac agt cat cta ggg cct 1298
 Gly Phe Ser Glu Glu Trp Glu Ala Gln Arg Asp Ser His Leu Gly Pro
 335 340 345 350
 cat cgc tct aca cct gag tca cga gct gct gtc cag gaa ctt tcc agc 1346
 His Arg Ser Thr Pro Glu Ser Arg Ala Ala Val Gln Glu Leu Ser Ser
 355 360 365
 agt atc ctc gct ggt gaa gac cca gag gaa agg gga gta aaa ctt gga 1394
 Ser Ile Leu Ala Gly Glu Asp Pro Glu Glu Arg Gly Val Lys Leu Gly
 370 375 380
 ttg gga gat ttc att ttc tac agt gtt ctg gtt ggt aaa gcc tca gca 1442
 Leu Gly Asp Phe Ile Phe Tyr Ser Val Leu Val Gly Lys Ala Ser Ala
 385 390 395
 aca gcc agt gga gac tgg aac aca acc ata gcc tgt ttc gta gcc ata 1490
 Thr Ala Ser Gly Asp Trp Asn Thr Thr Ile Ala Cys Phe Val Ala Ile
 400 405 410
 tta att ggt ttg tgc ctt aca tta tta ctc ctt gcc att ttc aag aaa 1538
 Leu Ile Gly Leu Cys Leu Thr Leu Leu Leu Leu Ala Ile Phe Lys Lys
 415 420 425 430
 gca ttg cca gct ctt cca atc tcc atc acc ttt ggg ctt gtt ttc tac 1586
 Ala Leu Pro Ala Leu Pro Ile Ser Ile Thr Phe Gly Leu Val Phe Tyr
 435 440 445
 ttt gcc aca gat tat ctt gta cag cct ttt atg gac caa tta gca ttc 1634
 Phe Ala Thr Asp Tyr Leu Val Gln Pro Phe Met Asp Gln Leu Ala Phe
 450 455 460
 cat caa ttt tat atc tag catatttgcg gttagaatcc catggatggt 1682
 His Gln Phe Tyr Ile
 465
 tcttctttga ctataaccaa atctggggag gacaaagggtg attttcctgt gtccacatct 1742
 aacaaagtca agattcccg ctggactttt gcagcttccct tccaagtctt cctgaccacc 1802
 ttgcactatt ggactttgga aggaggtgcc tatagaaaac gattttgaac atacttcac 1862
 gcagtggaact gtgtccctcg gtgcagaaac taccagattt gagggacgag gtcaaggaga 1922
 tatgataggc ccggaagtgt ctgtgccccca tcagcagctt gacgcgtggt cacaggacga 1982
 tttcactgac actgcgaact ctgaggacta ccggttacca agagggttagg tgaagtgggt 2042
 taaaccaaac ggaactcttc atcttaaaact acacgttgaa aatcaaccca ataattctgt 2102
 attaactgaa ttctgaactt ttcaggagggt actgtgagga agagcaggca ccagcagcag 2162
 aatggggaat ggagaggtgg gcagggggtc cagcttccct ttgatttttt gctgcagact 2222
 catccttttt aaatgagact tgttttcccc tctctttgag tcaagtcaaa tagtagatt 2282
 gcctttggca attcttcttc tcaagcactg acactcatta ccgtctgtga ttgccatttc 2342
 ttcccaaggc cagtctgaac ctgaggttgc tttatcctaa aagttttaac ctgaggttcc 2402

aaattcagta aattttggaa acagtacagc tattttctcat caattctcta tcattgtttaa 2462
 gtcaaatgtg gattttccac caaattctga atttgtagac atactgttac gctcacttgc 2522
 cccagatgc ctctctgtc ctctctctc tctccacac aagcagtctt tttctacagc 2582
 cagtaaggca gctctgtctt ggtagcagat ggtccattta ttctagggtc ttactctttg 2642
 tatgatgaaa agaattgtgt atgaatcggg gctgtcagcc ctgctgtcag acctctctcc 2702
 acagcaaatg agatgtatgc ccaaagcggg agaattaaag aagagtaaaa tggctgttga 2762
 agc 2765

<210> 5

<211> 448

<212> PRT

<213> Homo sapiens

<400> 5

Met	Leu	Thr	Phe	Met	Ala	Ser	Asp	Ser	Glu	Glu	Glu	Val	Cys	Asp	Glu	1	5	10	15
Arg	Thr	Ser	Leu	Met	Ser	Ala	Glu	Ser	Pro	Thr	Pro	Arg	Ser	Cys	Gln	20	25	30	
Glu	Gly	Arg	Gln	Gly	Pro	Glu	Asp	Gly	Glu	Asn	Thr	Ala	Gln	Trp	Arg	35	40	45	
Ser	Gln	Glu	Asn	Glu	Glu	Asp	Gly	Glu	Glu	Asp	Pro	Asp	Arg	Tyr	Val	50	55	60	
Cys	Ser	Gly	Val	Pro	Gly	Arg	Pro	Pro	Gly	Leu	Glu	Glu	Leu	Thr		65	70	75	80
Leu	Lys	Tyr	Gly	Ala	Lys	His	Val	Ile	Met	Leu	Phe	Val	Pro	Val	Thr	85	90	95	
Leu	Cys	Met	Ile	Val	Val	Val	Ala	Thr	Ile	Lys	Ser	Val	Arg	Phe	Tyr	100	105	110	
Thr	Glu	Lys	Asn	Gly	Gln	Leu	Ile	Tyr	Thr	Thr	Phe	Thr	Glu	Asp	Thr	115	120	125	
Pro	Ser	Val	Gly	Gln	Arg	Leu	Leu	Asn	Ser	Val	Leu	Asn	Thr	Leu	Ile	130	135	140	
Met	Ile	Ser	Val	Ile	Val	Met	Thr	Ile	Phe	Leu	Val	Val	Leu	Tyr		145	150	155	160
Lys	Tyr	Arg	Cys	Tyr	Lys	Phe	Ile	His	Gly	Trp	Leu	Ile	Met	Ser	Ser	165	170	175	
Leu	Met	Leu	Leu	Phe	Leu	Phe	Thr	Tyr	Ile	Tyr	Leu	Gly	Glu	Val	Leu	180	185	190	
Lys	Thr	Tyr	Asn	Val	Ala	Met	Asp	Tyr	Pro	Thr	Leu	Leu	Leu	Thr	Val	195	200	205	
Trp	Asn	Phe	Gly	Ala	Val	Gly	Met	Val	Cys	Ile	His	Trp	Lys	Gly	Pro	210	215	220	
Leu	Val	Leu	Gln	Gln	Ala	Tyr	Leu	Ile	Met	Ile	Ser	Ala	Leu	Met	Ala	225	230	235	240
Leu	Val	Phe	Ile	Lys	Tyr	Leu	Pro	Glu	Trp	Ser	Ala	Trp	Val	Ile	Leu	245	250	255	
Gly	Ala	Ile	Ser	Val	Tyr	Asp	Leu	Val	Ala	Val	Leu	Cys	Pro	Lys	Gly	260	265	270	
Pro	Leu	Arg	Met	Leu	Val	Glu	Thr	Ala	Gln	Glu	Arg	Asn	Glu	Pro	Ile	275	280	285	
Phe	Pro	Ala	Leu	Ile	Tyr	Ser	Ser	Ala	Met	Val	Trp	Thr	Val	Gly	Met	290	295	300	
Ala	Lys	Leu	Asp	Pro	Ser	Ser	Gln	Gly	Ala	Leu	Gln	Leu	Pro	Tyr	Asp	305	310	315	320
Pro	Glu	Met	Glu	Glu	Asp	Ser	Tyr	Asp	Ser	Phe	Gly	Glu	Pro	Ser	Tyr	325	330	335	
Pro	Glu	Val	Phe	Glu	Pro	Pro	Leu	Thr	Gly	Tyr	Pro	Gly	Glu	Glu	Leu				

340 345 350
 Glu Glu Glu Glu Glu Arg Gly Val Lys Leu Gly Leu Gly Asp Phe Ile
 355 360 365
 Phe Tyr Ser Val Leu Val Gly Lys Ala Ala Ala Thr Gly Ser Gly Asp
 370 375 380
 Trp Asn Thr Thr Leu Ala Cys Phe Val Ala Ile Leu Ile Gly Leu Cys
 385 390 395 400
 Leu Thr Leu Leu Leu Ala Val Phe Lys Lys Ala Leu Pro Ala Leu
 405 410 415
 Pro Ile Ser Ile Thr Phe Gly Leu Ile Phe Tyr Phe Ser Thr Asp Asn
 420 425 430
 Leu Val Arg Pro Phe Met Asp Thr Leu Ala Ser His Gln Leu Tyr Ile
 435 440 445

<210> 6

<211> 2236

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (368) (1714)

<400> 6

cgagcggcgg cggagcaggc atttccagca gtgaggagac agccagaagc aagctattgg 60
 agctgaagga acctgagaca gaagctagtc cccctctga attttactga tgaagaaact 120
 gaggccacag agctaaagtg acttttccca aggtcgccca gcgaggacgt gggacttctc 180
 agacgtcagg agagtgatgt gagggagctg tgtgaccata gaaagtgcg tgtaaaaac 240
 cagcgctgcc ctctttgaaa gccaggagc atcattcatt tagcctgctg agaagaagaa 300
 accaagtgtc cgggattcag acctctctgc ggccccaagt gttcgtggtg cttccagagg 360
 cagggtc atg ctc aca ttc atg gcc tct gac agc gag gaa gaa gtg tgt 409
 Met Leu Thr Phe Met Ala Ser Asp Ser Glu Glu Glu Val Cys
 1 5 10
 gat gag cgg acg tcc cta atg tcg gcc gag agc ccc acg ccg cgc tcc 457
 Asp Glu Arg Thr Ser Leu Met Ser Ala Glu Ser Pro Thr Pro Arg Ser
 15 20 25 30
 tgc cag gag ggc agg cag ggc cca gag gat gga gag aac act gcc cag 505
 Cys Gln Glu Gly Arg Gln Gly Pro Glu Asp Gly Glu Asn Thr Ala Gln
 35 40 45
 tgg aga agc cag gag aac gag gag gac ggt gag gag gac cct gac cgc 553
 Trp Arg Ser Gln Glu Asn Glu Glu Asp Gly Glu Glu Asp Pro Asp Arg
 50 55 60
 tat gtc tgt agt ggg gtt ccc ggg cgg ccg cca ggc ctg gag gaa gag 601
 Tyr Val Cys Ser Gly Val Pro Gly Arg Pro Pro Gly Leu Glu Glu Glu
 65 70 75
 ctg acc ctc aaa ttc gga gcg aag cac gtg atc atg ctg ttt gtg cct 649
 Leu Thr Leu Lys Tyr Gly Ala Lys His Val Ile Met Leu Phe Val Pro
 80 85 90
 gtc act ctg tgc atg atc gtg gtg gta gcc acc atc aag tct gtg cgc 697
 Val Thr Leu Cys Met Ile Val Val Val Ala Thr Ile Lys Ser Val Arg
 95 100 105 110
 ttc tac aca gag aag aat gga cag ctc atc tac acg aca ttc act gag 745
 Phe Tyr Thr Glu Lys Asn Gly Gln Leu Ile Tyr Thr Thr Phe Thr Glu
 115 120 125
 gac aca ccc tcg gtg ggc cag cgc ctc ctc aac tcc gtg ctg aac acc 793
 Asp Thr Pro Ser Val Gly Gln Arg Leu Leu Asn Ser Val Leu Asn Thr
 130 135 140
 ctc atc atg atc agc gtc atc gtg gtt atg acc atc ttc ttg gtg gtg 841

Leu Ile Met Ile Ser Val Ile Val Val Met Thr Ile Phe Leu Val Val
 145 150 155
 ctc tac aag tac cgc tgc tac aag ttc atc cat ggc tgg ttg atc atg 889
 Leu Tyr Lys Tyr Arg Cys Tyr Lys Phe Ile His Gly Trp Leu Ile Met
 160 165 170
 tct tca ctg atg ctg ctg ttc ctc ttc acc tat atc tac ctt ggg gaa 937
 Ser Ser Leu Met Leu Leu Phe Leu Phe Thr Tyr Ile Tyr Leu Gly Glu
 175 180 185 190
 gtg ctc aag acc tac aat gtg gcc atg gac tac ccc acc ctc ttg ctg 985
 Val Leu Lys Thr Tyr Asn Val Ala Met Asp Tyr Pro Thr Leu Leu Leu
 195 200 205
 act gtc tgg aac ttc ggg gca gtg ggc atg gtg tgc atc cac tgg aag 1033
 Thr Val Trp Asn Phe Gly Ala Val Gly Met Val Cys Ile His Trp Lys
 210 215 220
 ggc cct ctg gtg ctg cag cag gcc tac ctc atc atg atc agt gcg ctc 1081
 Gly Pro Leu Val Leu Gln Gln Ala Tyr Leu Ile Met Ile Ser Ala Leu
 225 230 235
 atg gcc cta gtg ttc atc aag tac ctc cca gag tgg tcc gcg tgg gtc 1129
 Met Ala Leu Val Phe Ile Lys Tyr Leu Pro Glu Trp Ser Ala Trp Val
 240 245 250
 atc ctg ggc gcc atc tct gtg tat gat ctc gtg gct gtg ctg tgt ccc 1177
 Ile Leu Gly Ala Ile Ser Val Tyr Asp Leu Val Ala Val Leu Cys Pro
 255 260 265 270
 aaa ggg cct ctg aga atg ctg gta gaa act gcc cag gag aga aat gag 1225
 Lys Gly Pro Leu Arg Met Leu Val Glu Thr Ala Gln Glu Arg Asn Glu
 275 280 285
 ccc ata ttc cct gcc ctg ata tac tca tct gcc atg gtg tgg acg gtt 1273
 Pro Ile Phe Pro Ala Leu Ile Tyr Ser Ser Ala Met Val Trp Thr Val
 290 295 300
 ggc atg gcg aag ctg gac ccc tcc tct cag ggt gcc ctc cag ctc ccc 1321
 Gly Met Ala Lys Leu Asp Pro Ser Ser Gln Gly Ala Leu Gln Leu Pro
 305 310 315
 tac gac ccg gag atg gaa gaa gac tcc tat gac agt ttt ggg gag cct 1369
 Tyr Asp Pro Glu Met Glu Glu Asp Ser Tyr Asp Ser Phe Gly Glu Pro
 320 325 330
 tca tac ccc gaa gtc ttt gag cct ccc ttg act ggc tac cca ggg gag 1417
 Ser Tyr Pro Glu Val Phe Glu Pro Pro Leu Thr Gly Tyr Pro Gly Glu
 335 340 345 350
 gag ctg gag gaa gag gag gaa agg ggc gtg aag ctt ggc ctc ggg gac 1465
 Glu Leu Glu Glu Glu Glu Glu Arg Gly Val Lys Leu Gly Leu Gly Asp
 355 360 365
 ttc atc ttc tac agt gtg ctg gtg ggc aag gcg gct gcc acg ggc agc 1513
 Phe Ile Phe Tyr Ser Val Leu Val Gly Lys Ala Ala Ala Thr Gly Ser
 370 375 380
 ggg gac tgg aat acc acg ctg gcc tgc ttc gtg gcc atc ctc att ggc 1561
 Gly Asp Trp Asn Thr Thr Leu Ala Cys Phe Val Ala Ile Leu Ile Gly
 385 390 395
 ttg tgt ctg acc ctc ctg ctg ctt gct gtg ttc aag aag gcg ctg ccc 1609
 Leu Cys Leu Thr Leu Leu Leu Leu Ala Val Phe Lys Lys Ala Leu Pro
 400 405 410
 gcc ctc ccc atc tcc atc acg ttc ggg ctc atc ttt tac ttc tcc acg 1657
 Ala Leu Pro Ile Ser Ile Thr Phe Gly Leu Ile Phe Tyr Phe Ser Thr
 415 420 425 430
 gac aac ctg gtg ccg ccg ttc atg gac acc ctg gcc tcc cat cag ctc 1705
 Asp Asn Leu Val Arg Pro Phe Met Asp Thr Leu Ala Ser His Gln Leu
 435 440 445
 tac atc tga gggacatggt gtgccacagg ctgcaagctg cagggaattt 1754
 Tyr Ile

```
tcattggatg cagttgtata gttttacact ctagtgccat atatttttaa gacttttctt 1814
tccttaaaaa ataaagtacg tgtttacttg gtgaggagga ggcagaacca gctctttggt 1874
gccagctgtt tcatcaccag actttggctc ccgctttggg gagcgccctg cttcacggac 1934
aggaagcaca gcaggtttat ccagatgaac tgagaaggctc agattagggc ggggagaaga 1994
gcatccggca tgagggctga gatgcgcaaa gagtgtgctc gggagtggcc cctggcacct 2054
gggtgctctg gctggagagg aaaagccagt tccctacgag gagtgttccc aatgctttgt 2114
ccatgatgtc cttgttattt tattgccttt agaaactgag tcctgttctt gttacggcag 2174
tcacactgct ggggaagtggc ttaatagtaa tatcaataaa tagatgagtc ctgttagaaa 2234
aa
```


(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 September 2001 (13.09.2001)

PCT

(10) International Publication Number
WO 01/67097 A3

- (51) International Patent Classification⁷: **G01N 33/68**
- (21) International Application Number: **PCT/GB01/01059**
- (22) International Filing Date: **9 March 2001 (09.03.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
0005895.8 10 March 2000 (10.03.2000) **GB**
- (71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED [GB/GB]**; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HALE, Richard, Stephen [GB/GB]**; GlaxoSmithKline plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **ROWLEY, Adele [GB/GB]**; GlaxoSmithKline plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **BLACKSTOCK, Walter [GB/GB]**; GlaxoSmithKline plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB).
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO** patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), **Eurasian** patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), **European** patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), **OAPI** patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
- (88) Date of publication of the international search report:
16 May 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **METHOD OF IDENTIFYING MODULATORS OF PRESENILIN**

(57) Abstract: A method of identifying a modulator of presenilin function, the method comprising: (i) providing (a) a polypeptide capable of binding a presenilin, which polypeptide comprises the amino acid sequence of SEQ ID NO: 1 or a functional variant thereof, or a fragment of either thereof which is capable of binding to presenilin; (b) a presenilin or a variant thereof or a fragment of either thereof capable of binding to a polypeptide which comprises the amino acid sequence of SEQ ID NO: 1; (c) a test substance under conditions that would permit binding of a polypeptide (a) to a presenilin (b) in the absence of the test substance; (ii) monitoring presenilin mediated activity; and (iii) determining thereby whether the test substance is a modulator of presenilin activity, a modulator identified by a method of the invention and use of a modulator identified by a method of the invention in the treatment of Alzheimer's disease.

WO 01/67097 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/01059

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, MEDLINE, SCISEARCH, CHEM ABS Data, BIOSIS, EMBASE, SEQUENCE SEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 35501 A (UNIV TORONTO) 15 July 1999 (1999-07-15) the whole document	1-10, 13-16
Y	KEIRSEBILCK A ET AL: "MOLECULAR CLONING OF THE HUMAN P120 CTN CATENIN GENE (CTNND1): EXPRESSION OF MULTIPLE ALTERNATIVELY SPLICED ISOFORMS" GENOMICS, ACADEMIC PRESS, SAN DIEGO, US, vol. 50, 1998, pages 129-146, XP000864646 ISSN: 0888-7543 cited in the application the whole document	1-10, 13-16

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

14 January 2002

Date of mailing of the international search report

29/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2260 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Niemann, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/01059

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>ANASTASIADIS PANOS Z ET AL: "The p120 catenin family: Complex roles in adhesion, signaling and cancer." JOURNAL OF CELL SCIENCE, vol. 113, no. 8, April 2000 (2000-04), pages 1319-1334, XP002187294 ISSN: 0021-9533</p>	

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 11-12

Present claims 11 and 12 relate to a compound or the use of that compound defined by reference to a desirable characteristic or property, namely being identifiable by the screening methods of claims 1-10.

The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for no such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the screening methods per se.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/01059

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9935501 A	15-07-1999	AU 1957299 A WO 9935501 A1	26-07-1999 15-07-1999

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.